

Male and female brains aren't equal when it comes to fat

October 16 2014

Researchers have found that male and female brains respond in remarkably different ways to high-fat meals. Those differences in the brain lead to greater inflammation and increased health risks in males that indulge on fatty foods in comparison to females, a new study in mice shows. The findings reported in the Cell Press journal *Cell Reports* on October 16th may help to explain observed differences in obesity outcomes between women and men – premenopausal women carrying extra weight fare better than men do – and suggest that dietary advice should be made more sex-specific.

"Our findings, for the first time, suggest that males and females respond to high-fat diets differently," said Deborah Clegg of the Cedar-Sinai Diabetes And Obesity Research Institute in Los Angeles. "The data would suggest that is probably 'ok' for females to occasionally have a high-fat meal, where it is not recommended for males.

"The way we treat patients and provide dietary and nutritional advice should be altered. We might be less concerned about an occasional hamburger for women, but for men, we might more strongly encourage avoidance, especially if they have pre-existing diseases such as heart disease or type 2 diabetes."

Earlier data from Clegg's team and others had suggested that [inflammation](#) in the brain is tied to overeating, blood sugar imbalances, and increased inflammation in other parts of the body, including fat tissue. Those effects can be triggered, in males in particular, by short-

term exposure to a high-fat diet.

The researchers say they were initially shocked to discover that male and female brains differ in their fatty acid composition. When they manipulated male mouse brains to have the fatty acid profile of females, they found that those animals were protected from the ill effects of a diet high in fat.

When males with average male brains entered an inflammatory state after eating diets high in fat, they also suffered from reduced cardiac function in a way that female animals in the study did not. Those sex differences in the brain's response to fat are related to differences between females and males in estrogen and estrogen receptor status.

Clegg says her team is now working out a strategy to confirm whether the findings in mice apply to people too. If they do, there will be some very immediate practical implications for what men and women should put on their plates.

"We have always had 'one size fits all' with respect to our nutritional information and our pharmaceutical approach," Clegg said. "Our data begin to suggest that sex should be factored in, and men should be more closely monitored for fat intake and inflammation than women."

More information: *Cell Reports* [www.cell.com/cell-reports/abstract/2211-1247\(14\)00808-0](http://www.cell.com/cell-reports/abstract/S2211-1247(14)00808-0)

Provided by Cell Press

Citation: Male and female brains aren't equal when it comes to fat (2014, October 16) retrieved 26 April 2024 from <https://medicalxpress.com/news/2014-10-male-female-brains-equal-fat.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.