

New optimal screening threshold for gestational diabetes in twin pregnancies

October 30 2014

A common complication, gestational diabetes affects approximately 6-7% of pregnant women. Currently, screening is done in two steps to help identify patients most at risk; however, the suggested levels for additional testing were based on singleton pregnancy data. Now investigators have analyzed data from twin pregnancies and have determined that the optimal first step cutoff for additional screening appears to be a blood sugar level equal to or greater than 135 mg/dL for women carrying twins. Their findings are published in the *American Journal of Obstetrics & Gynecology*.

Gestational diabetes mellitus (GDM) is believed to be caused by increased levels of human placental lactogen, estrogen, cortisol, prolactin, and progesterone. GDM is linked to several adverse birth outcomes such as preeclampsia, macrosomia, birth injury, and cesarean delivery. Because of the possible negative effects, screening for GDM is an important component of prenatal care.

Representing approximately 3.3% of all live births in the United States, twin pregnancies present unique medical challenges; however, most of the available data regarding GDM testing thresholds are based on singleton pregnancies. Since carrying twins amplifies the normal physical changes that accompany pregnancy, a team of researchers from the Icahn School of Medicine at Mount Sinai and the New York University School of Medicine set out to pinpoint the most effective screening cutoff for women pregnant with twins.



Currently, the American Congress of Obstetricians and Gynecologists advocates a two-step diagnostic approach. Women are screened between 24-28 weeks, using a nonfasting 1-hour 50-g glucose challenge test (GCT). If a patient exceeds the recommended cutoff (between \geq 130 mg/dL and \geq 140 mg/dL), additional testing is performed to diagnose or rule out GDM.

In this study, doctors analyzed data from 475 women pregnant with twins who were treated at their hospitals between 2005 and 2013. Specifically, they focused on three potential cutoff thresholds for the GCT: \geq 130 mg/dL, \geq 135 mg/dL, and \geq 140 mg/dL. The investigators found that the optimal cutoff for the 1-hour 50-g GCT screening was \geq 135 mg/dL. The data revealed that the 130 mg/dL cutoff did not show any additional predictive value, yet increased the test positive rate. The 140 mg/dL cutoff would have failed to identify 6.5% of patients with GDM.

"With the increasing prevalence of twin pregnancies, understanding how to better screen for GDM is important in this population because of the unique circumstances that may put this group at increased risk for development of insulin resistance, as well as their altered physiology, which would limit the ability to extrapolate from data derived from singleton pregnancies," says investigator Nathan S. Fox, MD, Associate Clinical Professor, Obstetrics, Gynecology and Reproductive Science, Icahn School of Medicine at Mount Sinai. "We found that the optimal GCT cutoff appears to be ≥ 135 mg/dL. This cutoff yielded 100% sensitivity with only a 28.6% test positive rate."

While early detection and screening are essential for effective treatment and control of GDM, unnecessary follow-up testing can cause mental stress for the mother and result in additional out-of-pocket expenditures. "As opposed to a cutoff of \geq 130 mg/dL, a cutoff of \geq 135 mg/dL could potentially decrease cost and anxiety, without decreasing the sensitivity



of the test," explains Dr. Fox.

For now, this study represents an important first step in addressing the distinctive differences between GDM in twin and singleton pregnancies, but more research is needed to fully develop guidelines to better serve mothers carrying twins. "There is much more to learn regarding GDM in twin pregnancies that could be addressed with future prospective studies," concludes Dr. Fox. "More research is needed in twin pregnancies to confirm this finding in other populations, as well as to establish the optimal GDM screening and treatment paradigm in twin pregnancies."

More information: "Screening approach for gestational diabetes in twin pregnancies," by Andrei Rebarber, MD; Cara Dolin, MD; Jessica C. Fields, BA; Daniel H. Saltzman, MD; Chad K. Klauser, MD; Simi Gupta, MD; Nathan S. Fox, MD (DOI: <u>dx.doi.org/10.1016/j.ajog.2014.08.030</u>), online in advance of the *American Journal of Obstetrics & Gynecology*, Volume 211, Issue 6 (December 2014)

Provided by Elsevier

Citation: New optimal screening threshold for gestational diabetes in twin pregnancies (2014, October 30) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2014-10-optimal-screening-threshold-gestational-diabetes.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.