

Pharmacoinvasive STEMI strategy best for smokers, nonsmokers

October 27 2014



(HealthDay)—For patients with ST-segment elevation myocardial infarction, a pharmacoinvasive strategy after fibrinolysis is beneficial for smokers and nonsmokers, according to a study published in the Oct. 1 issue of *The American Journal of Cardiology*.

Nigel S. Tan, M.D., from St. Michael's Hospital in Toronto, and colleagues examined whether the optimal treatment strategy after fibrinolysis differs based on <u>smoking status</u>. Data were collected from patients with ST-segment elevation <u>myocardial infarction</u> who were randomized to a routine early invasive (pharmacoinvasive) versus a standard (early transfer only for rescue <u>percutaneous coronary intervention</u> or delayed angiography) strategy after fibrinolysis. The interaction between smoking status and treatment strategy was examined.



The researchers found that after multivariate adjustment, smoking status was not a significant predictor of either primary or secondary end points. Compared with standard therapy, pharmacoinvasive management reduced the primary end point in smokers (7.7 versus 13.6 percent; P = 0.04) and in nonsmokers (13.1 versus 19.7 percent; P = 0.03). The effect of treatment was not modified by smoking status for any measured outcomes (P > 0.10 for all).

"The beneficial treatment effect of a pharmacoinvasive strategy is consistent in smokers and <u>nonsmokers</u>," the authors write.

The trial was partially funded by a grant from Roche; coronary stents were provided free of charge by Abbott. Several authors disclosed financial ties to the pharmaceutical industry.

More information: Abstract

Full Text (subscription or payment may be required)

Copyright © 2014 HealthDay. All rights reserved.

Citation: Pharmacoinvasive STEMI strategy best for smokers, nonsmokers (2014, October 27) retrieved 16 June 2024 from https://medicalxpress.com/news/2014-10-pharmacoinvasive-stemi-strategy-smokers-nonsmokers.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.