

Teens playing high-contact sports at risk for using drugs, alcohol

October 22 2014, by Jared Wadley

(Medical Xpress)—Teens who play sports like football, wrestling, hockey or lacrosse are more likely to drink alcohol or smoke cigarettes or marijuana than student athletes who play noncontact sports, according to a new University of Michigan study.

The findings show that participation in high-contact sports is associated with substance use during the past 30 days. Meanwhile, participating in noncontact sports, such as tennis, swimming, gymnastics and track, lessened the likelihood of substance use in the past month.

"Competitive [sports participation](#) can either inhibit or amplify substance use. It just depends upon which type of sport adolescents are involved with," said Philip Veliz, assistant research professor at the Institute for Research on Women and Gender.

Veliz, the study's lead author, collaborated with colleagues Carol Boyd, the Deborah J. Oakley Collegiate Professor of Nursing and a professor of women's studies, and Sean McCabe, research professor at the Institute for Research on Women and Gender.

Contact sport participants view their body as an instrument that can be easily gambled with, even if it means permanent damage, Veliz said. Conversely, sports that involve minimal to no contact are valorized for their sustainability for participants throughout life.

"It is also assumed that youth who are involved with noncontact sports

will be the least likely to engage in substance use due to the emphasis placed on fostering a strategic orientation to maintain a level of fitness for both competition and future longevity," he said.

The researchers used data from Monitoring the Future, which surveys 8th-, 10th- and 12th-grade students on a range of different topics like substance use, academic performance and competitive sports participation. The national sample included more than 21,000 teens who were asked about substance and illicit drug use during a 30-day period.

The questions were grouped into three categories: high-contact sports (involve continual violent contact that sanctions hitting, knocking, wrestling or tackling opponents to the ground); semicontact sports (have sporadic violent contact that may be officially sanctioned within the rules of the sport, such as baseball, basketball, field hockey or soccer); and noncontact sports (have no contact between participants due to either official rules or the structure of the playing field).

The study also indicated that respondents in competitive [sports](#) had higher odds of indicating early onset of getting drunk between 4th and 8th grades when compared to their nonparticipating peers.

Veliz said the findings, unfortunately, cast doubt on the belief that participation in any type of competitive sport deters adolescents from engaging in risky behaviors like substance use.

More information: "Competitive Sport Involvement and Substance Use among Adolescents: A Nationwide Study." Philip Todd Veliz, Carol J. Boyd, Sean Esteban McCabe. *Substance Use & Misuse* . [DOI: 10.3109/10826084.2014.962049](https://doi.org/10.3109/10826084.2014.962049)

Provided by University of Michigan

Citation: Teens playing high-contact sports at risk for using drugs, alcohol (2014, October 22)
retrieved 23 April 2024 from

<https://medicalxpress.com/news/2014-10-teens-high-contact-sports-drugs-alcohol.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.