

Young adults aren't too young to be at risk for a stroke

October 30 2014, by Kim Irwin



Jennifer Reilly was 28 years old when she began experiencing some odd symptoms. Half of her left hand went numb. She could move her fingers, but she couldn't feel anything on the outer part of that hand.

Because she was so young, Reilly assumed it was nothing. She was wrong.

She was having a [stroke](#).

The numbness recurred off and on for several days. Then one night she had a piercing headache, an uncommon event for her. She mentioned her

[symptoms](#) to a work colleague, who urged her to see a doctor.

"I didn't know I was having a stroke at the time," said Reilly, now 35. "I just assumed I was a healthy, normal 28-year-old."

Reilly saw several doctors and finally landed at UCLA in the neurology department, where she saw Dr. David Liebeskind, professor of neurology and director of outpatient stroke and neurovascular programs. After a battery of tests, Reilly was diagnosed with Moyamoya disease—a rare, progressive cerebrovascular disorder caused by blocked arteries at the base of the brain, which cuts off blood flow. One of the first symptoms of Moyamoya is recurrent [transient ischemic attacks](#), or TIAs, commonly referred to as "mini-strokes," which are exactly what Reilly was experiencing.

She had no idea the danger she was in, said Liebeskind, who is also director of the neurovascular imaging research core at UCLA.

"The worst and a very likely possibility is that she would have had a significant stroke," he said, one that could have been extremely debilitating or even fatal.

Reilly was referred to Dr. Neil Martin, chair of neurosurgery at Ronald Reagan UCLA Medical Center and head of the neurovascular surgery section.

"He told me I needed to be admitted into the hospital immediately, and that I would be in surgery within 24 hours," she said. "It was such a shock. I can't fathom what my parents must have been going through. We had a few tears, and they stayed very strong for me."

Dr. Nestor Gonzalez, an associate professor of neurosurgery and radiology, performed two nine-hour brain surgeries on Reilly to reroute

arteries that normally bring blood to the scalp to instead supply blood to her brain. Over time, the arteries formed new branches and restored blood supply to the brain.

Gonzalez highlighted the importance of Reilly acting quickly after her symptoms began.

"The good thing is that she sought attention when she started having those symptoms and we were able to diagnose her condition very early," he said. "We were able to avert a stroke that could have devastated her life."

It's a common misconception that strokes only affect older people, Gonzalez said.

"A stroke is not a disease limited to elder individuals," he said. "It is a condition that can affect younger individuals, even kids. The moment someone feels any of the symptoms of a stroke, they need to seek immediate emergency attention."

Liebeskind said the most common signs of stroke include sudden onset of weakness, sudden onset of numbness, difficulty speaking, difficulty seeing and difficulty interacting with other individuals.

It's been nearly six years since her mini strokes and surgeries and today Reilly is embracing life. She got married and will celebrate her third wedding anniversary next June.

"Since my recovery I have an absolutely newfound perspective on life," she said. "I really try to stop and appreciate that I was given a second chance."

Provided by University of California, Los Angeles

Citation: Young adults aren't too young to be at risk for a stroke (2014, October 30) retrieved 18 April 2024 from <https://medicalxpress.com/news/2014-10-young-adults.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.