

# HIV-infected adults diagnosed with age-related diseases at similar ages as uninfected

November 11 2014

---

New research from the Johns Hopkins Bloomberg School of Public Health suggests that HIV-infected adults are at a higher risk for developing heart attacks, kidney failure and cancer. But, contrary to what many had believed, the researchers say these illnesses are occurring at similar ages as adults who are not infected with HIV.

The findings appeared online last month in the journal *Clinical Infectious Diseases*.

Researchers say these findings can help reassure HIV-infected patients and their [health care providers](#).

"We did not find conclusive evidence to suggest that screening for these diseases should occur at younger ages in HIV-infected compared with uninfected adults," says Keri N. Althoff, PhD, MPH, an assistant professor in the Department of Epidemiology at the Bloomberg School and lead author of the study.

Using data from 98,687 HIV-infected and uninfected adults in the Veterans Aging Cohort Study from April 1, 2003, to Dec. 31, 2010, the researchers were able to compare the ages at which each group was diagnosed with heart attacks, [kidney failure](#) and cancers unrelated to AIDS. Althoff and her colleagues found that HIV-infected adults were diagnosed with kidney failure, on average, when they were six months younger than the group without HIV. There were no statistically significant age differences at diagnosis between infected and uninfected

adults for heart attacks or cancers unrelated to AIDS.

The study also compared the risk of these age-related diseases in HIV-infected and uninfected adults. Similar to previous studies, HIV-infected adults appear to be at an increased risk of these diseases, but the magnitude of the increase in risk was smaller than has been reported in many of the previous studies, likely due to this study's ability to adjust for factors such as race, gender, obesity, smoking, diabetes, and depression, among others, influencing the risk in both HIV-infected and uninfected adults.

In the early years of the AIDS epidemic, HIV was a death sentence. But in recent years antiretroviral therapy (ART) has been proven highly effective at suppressing HIV viral replication and increasing longevity for people with HIV. A study published in December 2013 in the North American AIDS Cohort Collaboration on Research and Design (NA-ACCORD) shows that the life expectancy among HIV-infected adults is only a few years shy of the U.S. general population.

While HIV-infected people are living longer, there has been a contentious debate in medical circles about whether they are undergoing an accelerated aging process. People living with HIV have demanded a more rigorous study of whether and why this may be happening.

Just because the data show that the onset of these three diseases is similar between those with and without HIV doesn't mean those with HIV are just as healthy, Althoff cautions.

"Many HIV-infected adults feel older than their age," she says. "Our study looks at three important age-related diseases and we find no meaningful difference in age at diagnosis. But this isn't the whole story. We need more research to figure out what is going on."

**More information:** "Comparison of risk and age at diagnosis of myocardial infarction, end-stage renal disease, and non-AIDS-defining cancer in HIV-infected vs. uninfected adults" *Clinical Infectious Diseases*, 2014.

Provided by Johns Hopkins University Bloomberg School of Public Health

Citation: HIV-infected adults diagnosed with age-related diseases at similar ages as uninfected (2014, November 11) retrieved 23 April 2024 from <https://medicalxpress.com/news/2014-11-hiv-infected-adults-age-related-diseases-similar.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.