

Periodontal disease linked to increased risk of kidney disease

November 14 2014

In a study of blacks with normal kidney function, those with severe periodontal disease developed chronic kidney disease (CKD) at 4 times the rate of those without severe periodontal disease. The study that will be presented at ASN Kidney Week 2014 November 11–16 at the Pennsylvania Convention Center in Philadelphia, PA.

Periodontal disease is a chronic bacterial infection of the [oral cavity](#), and it disproportionately affects African Americans. It's also been implicated as a potential risk factor for CKD. To investigate this potential link, researchers led by Vanessa Grubbs, MD (University of California, San Francisco) analyzed 699 African American adults who underwent complete dental examinations.

During an average follow-up of 4.8 years, there were 21 (3.0%) new CKD cases. Participants with severe periodontal disease had a 4.2-fold greater incidence of CKD after adjusting for various factors (age, sex, diabetes, hypertension, smoking, and income) compared with those without severe periodontal disease.

"Because periodontal disease is common and can be prevented and treated, targeting it may be an important path towards reducing existing racial and ethnic disparities in chronic and end-stage kidney disease," said Dr. Grubbs.

More information: Study: "The Association Between Periodontal Disease and Kidney Function Decline in African Americans: the Jackson

Heart Study" (Abstract FR-OR916)

Provided by American Society of Nephrology

Citation: Periodontal disease linked to increased risk of kidney disease (2014, November 14)
retrieved 9 April 2024 from

<https://medicalxpress.com/news/2014-11-periodontal-disease-linked-kidney.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--