


ASH: Reducing factor XI cuts clots after knee arthroplasty

December 8 2014



(HealthDay)—Reducing factor XI levels with a second-generation antisense oligonucleotide (FXI-ASO) is effective for preventing venous thromboembolism after total knee arthroplasty, according to a study published online Dec. 7 in the *New England Journal of Medicine*. The findings were released to coincide with the annual meeting of the American Society of Hematology, held from Dec. 6 to 9 in San Francisco.

Harry R. Büller, M.D., from the University of Amsterdam, and colleagues conducted an open-label parallel-group study involving 300 patients undergoing elective primary unilateral [knee arthroplasty](#). Participants were randomized to receive one of two doses of FXI-ASO (200 or 300 mg) or 40 mg of enoxaparin once daily.

The researchers found that around the time of surgery, the mean factor XI levels were 0.38, 0.20, and 0.93 units per milliliter in the 200-mg FXI-ASO, 300-mg FXI-ASO, and enoxaparin groups, respectively. The primary efficacy outcome of incidence of [venous thromboembolism](#) occurred in 27 percent of those who received the 200-mg dose of FXI-ASO, 4 percent of those who received the 300-mg dose of FXI-ASO, and 30 percent of those who received enoxaparin. Compared with enoxaparin, the 200-mg regimen was noninferior, and the 300-mg regimen was superior (P

"These findings support the concept that with the use of strategies targeting factor XI, thrombosis and hemostasis can be dissociated," the authors write.

The study was funded by Isis Pharmaceuticals, the manufacturer of FXI-ASO.

More information: [Abstract](#)

[Full Text](#)

[Editorial](#)

[More Information](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

Citation: ASH: Reducing factor XI cuts clots after knee arthroplasty (2014, December 8) retrieved 25 April 2024 from <https://medicalxpress.com/news/2014-12-ash-factor-xi-clots-knee.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.