

Basophil reactivity to allergens varies by time of day

December 9 2014



Image courtesy of Blausen Medical

(HealthDay)—Circadian variations in gene expression may contribute to temporal variations in the symptoms of seasonal allergic rhinitis, according to research published online Dec. 2 in *Allergy*.

Noriko Ando, M.D., of the University of Yamanashi in Japan, and colleagues examined allergen-specific basophil reactivity according to circadian clock activity in murine models.

The researchers found that allergen-induced surface CD203c expression on basophils in seasonal <u>allergic rhinitis</u>, in response to Japanese cedar pollen, was dependent on time of day according to temporal variations of canonical circadian clock <u>gene expression</u>. Bone marrow-derived basophils generated from wild-type mice showed variation in immunoglobulin E-mediated interleukin 4 and histamine production that



was dependent on time of day. This pattern was not observed in bone marrow-derived basophils generated from *Clock*-mutated mice.

"Therefore, allergen-specific basophil reactivity shows daily variations depending on the <u>circadian clock</u> activity in basophils, which could partly explain temporal symptomatic variations in allergic rhinitis," the authors write.

More information: <u>Abstract</u> <u>Full Text</u>

Copyright © 2014 HealthDay. All rights reserved.

Citation: Basophil reactivity to allergens varies by time of day (2014, December 9) retrieved 27 April 2024 from https://medicalxpress.com/news/2014-12-basophil-reactivity-allergens-varies-day.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.