

Brain scans link frontal abnormalities to suicidal behaviors in adolescents, young adults

December 9 2014

Abnormalities in the prefrontal cortex and related brain areas are observed in adolescents who have attempted suicide, according to a report today at the American College of Neuropsychopharmacology annual meeting in Phoenix Arizona. The study suggests that deficits in frontal systems may be associated with risk for suicide attempts in youths with mood disorders.

Most [suicide attempts](#) occur in the context of [mood disorders](#), including [bipolar disorder](#) and [major depressive disorder](#). Bipolar disorder has a prevalence of 3-4% in the U.S. population, and 25-50% of those affected attempt suicide; 15-20 % of individuals with the disorder die as a result. Early intervention is needed as suicidal behavior often first presents in adolescence. The development of new interventions, however, requires a better characterization of how features of brain structure and function are linked to the development of suicidal behaviors.

Using magnetic resonance imaging (MRI), Hilary Blumberg and colleagues at the Yale School of Medicine examined brain structure and function in adolescents and young adults, 14 - 25 years of age. Sixty-eight participants with bipolar disorder, of whom 26 attempted suicide, were compared with 45 healthy volunteers matched for age and gender. The investigators found that, compared with bipolar patients who did not attempt suicide and healthy control subjects, the participants who

attempted suicide showed less integrity of white matter in key frontal brain systems, including the uncinate fasciculus, a fiber tract that connects the frontal lobe with key [brain areas](#) that control emotion, motivation, and memory. What was remarkable was that the deficits in the structural connections were linked to weaker connections between the [prefrontal cortex](#) and amygdala, suggesting that the white matter abnormalities disrupt the ability of these system components to work together. Further, there were links between the circuitry deficits and suicidal ideation, the number of suicide attempts and the relative lethality of those prior suicide attempts.

This work presents an important first step in understanding the neurobiology of how suicidal thoughts and behaviors are generated and may facilitate earlier identification of individuals at risk and development of targeted interventions to stop suicide.

Provided by American College of Neuropsychopharmacology

Citation: Brain scans link frontal abnormalities to suicidal behaviors in adolescents, young adults (2014, December 9) retrieved 6 May 2024 from <https://medicalxpress.com/news/2014-12-brain-scans-link-frontal-abnormalities.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--