

Men in recovery from Ebola should wear condoms for at least three months

December 16 2014



Image: Wikipedia

A new article reports that despite a clear lack of research on male survivors of Ebola, the current recommended practice of waiting at least three months after recovery to have unprotected sex should be upheld. This study was published today in *Reproductive Sciences*.

"Our exercise demonstrated that the current recommendations to prevent the sexual spread of Ebola are based on one mere observation," the researchers wrote. "Despite the evident need to conduct more research, for now, health care professionals should strongly recommend sexual abstinence or condom-protected encounters for at least three months."



Drs. Walter Cardona-Maya, Paula Velilla Hernandez, and Daniel Henao studied research from 1977 to 2007 on men in the convalescent phase of the Ebola virus (i.e. symptoms were fading and they were in recovery). They found only four studies that report male survivors of Ebola who donated convalescent semen. For these men, the virus persisted in their semen for an average of 66.6 days, and in one case, it persisted for 91 days.

The researchers continued, "The current Ebola Virus Disease outbreak is the longest and largest we have ever seen. Nonetheless, our results clearly demonstrate how much we ignore about it. Larger studies are needed to establish the social, clinical and biological determinants of this neglected disease."

Provided by SAGE Publications

Citation: Men in recovery from Ebola should wear condoms for at least three months (2014, December 16) retrieved 24 April 2024 from https://medicalxpress.com/news/2014-12-men-recovery-ebola-condoms-months.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.