

# Four reasons I won't have a prostate cancer blood test

December 11 2014, by Ian Haines

---



For many men, the down sides of PSA testing outweigh the benefits. Credit: Gerald Streiter/Flickr, CC BY-NC

Cancer Council Australia and the Prostate Cancer Foundation of Australia yesterday released new draft guidelines to help GPs counsel men who ask about prostate cancer tests. They advise GPs to explain the pros and cons of testing and, if the man wants to proceed, to give him a prostate specific antigen (PSA) blood test every two years between the

ages of 50 to 69.

Over the past few decades public health messages have drummed into us that early detection and treatment of diseases are key to good outcomes. Add to this the celebrity testimonials for prostate tests and non-celebrities who tell us their PSA test "saved their life" and it's easy to see why men think the tests are beneficial.

While some prostate cancers are harmful and require treatment, many are not. So the prevailing wisdom – that [early detection](#) and treatment is best – doesn't necessarily apply. At least 70% of men over 70 have [prostate cancer detected in autopsies](#), and only [3% of men](#) die because of prostate cancer.

I'm a 60-year-old male oncologist who has practised full-time for 36 years. Having "skin in the game" I've followed the testing debate closely since PSA was first used for screening in the early 1990s. Based on the evidence, if I was asymptomatic, I would not choose to have PSA test. Here's why.

## 1. PSA is a poor testing tool

Prostate specific antigen (PSA) is an enzyme secreted in large amounts by normal as well as cancerous prostate cells. Only small amounts of PSA leak into circulation from a normal prostate, but this increases with any prostatic disease, benign or malignant.

An elevated PSA very often does not indicate cancer. Just [one in four men](#) with a positive PSA test will have prostate cancer.

PSA tests also miss many cancers. A [2003 study](#) found that 21% of men who had a "normal" PSA of 2.6 to 3.9 at the end of a seven-year study did, in fact, have prostate cancer. Of the men with a PSA of 2.5 or less,

15% had cancer.

## 2. Prostate cancer isn't like other cancers

The point of a cancer screening test is that it can reliably detect lesions that, if removed, will reduce the chances of that patient later developing a life-threatening cancer.

This is certainly true for polyps and bowel cancer screening. It is also powerfully true for precancerous lesions of the cervix and cervical cancer. It is somewhat true for mammography and precancerous or early invasive breast lesions.

It is not at all true for prostate cancer screening. In the majority of cases, prostate cancer behaves more like an indolent condition and does not pose any threat to the patient's natural life span. This proportion of men [dying of other causes](#) continues to increase in the PSA era.

The only prostate screening study showing an [advantage for screening](#) has [very serious flaws](#), which have also been noted by the head of the American Cancer Society [Professor Otis Brawley](#) and [Professor Richard Ablin](#), who discovered the [prostate specific antigen](#) in the 1970s.

## 3. Surgery won't always cure you

The [only study](#) comparing radical surgery with *no treatment* found equivalent outcomes.

It concluded:

Among men with localized prostate cancer detected during the early era of PSA testing, radical prostatectomy did not significantly reduce all-

cause or prostate-cancer mortality, as compared with observation, through at least 12 years of follow-up. Absolute differences were less than three percentage points.

Radical surgery "cures" six in seven cases of prostate cancer. But this does not remove the uncertainty and doubt for men with prostate cancer, as most of the six in seven who are "cured" did not require treatment and most of the one in seven with dangerous cancers requiring cure will not be helped by the treatment.

#### **4. Detection and treatment comes with side effects**

[One in six men](#) will be diagnosed with prostate cancer during his lifetime. Their lives will be profoundly changed by this cancer diagnosis, whether or not they proceed with treatment.

I do not want the anxiety, depression and relationship changes that follow diagnosis, radical surgery, active surveillance or any regular monitoring.

I do not want to be impotent, which is very likely after radical treatment, or have urinary incontinence.

Even before treatment commences, after an abnormal PSA result, [men](#) are referred for a prostate biopsy: a surgical procedure that, even though it can indicate cancer, cannot give reliable information about how that cancer will behave.

I do not want the 1-2% risk of life-threatening [infections](#) caused by prostate biopsies.

#### **Bottom line**

I'm happy to take active positive steps to improve my health where it is proven that the benefits of the intervention outweigh the costs. But I'm not prepared to have my life ruled by a regular [blood test](#) like PSA that has no advantage.

*This story is published courtesy of [The Conversation](#) (under Creative Commons-Attribution/No derivatives).*

Source: The Conversation

Citation: Four reasons I won't have a prostate cancer blood test (2014, December 11) retrieved 3 May 2024 from <https://medicalxpress.com/news/2014-12-wont-prostate-cancer-blood.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.