

Brief observations miss autism symptoms in young children

January 12 2015



Terisa Gabrielsen and granddaughter Adelaide Burton interacting. A child pretending or engaging in make-believe play is one of the items on the M-CHAT-R checklist.

(Medical Xpress)—As the rate of children with autism in the U.S. continues to grow, a new study published today in top-ranked journal *Pediatrics* shows that medical professionals can't rely solely on their clinical judgment to detect autism risk.

The study finds that 10-20-minute observations, such as a pediatric exam, don't provide enough information about symptoms associated with autism. Within that short window, many [children](#) with autism display mostly typical behavior and may fail to receive a referral for further autism testing, even if a few autism symptoms are noticed.

"One of the biggest problems with early identification of autism is that many children aren't identified until they reach the school system," said the study's lead author and BYU assistant professor Terisa Gabrielsen. "This means that they have missed out on some prime years for intervention that can change a child's outcome."

Last year the CDC released a report stating that 1 in 68 U.S. children has an [autism spectrum disorder](#) (ASD). That's a 30 percent increase from 1 in 88 from two years earlier. These increases mean an increased demand for autism referral decisions

The study in *Pediatrics* looked at children 15-33 months old, with autism experts analyzing 10-minute videos of the children's behaviors during evaluation in a clinical setting. Children with autism, speech delays and typical children were included. The researchers wanted to document the ratio of typical behaviors vs. atypical behaviors exhibited and the corresponding referral decisions based on the observations. They found that within the brief timeframe of 10 minutes, children with autism exhibited much more typical behavior than atypical behavior overall, making it easy for clinicians to miss detecting autism risk. In the study, even the experts who reviewed the videos missed referrals for 39 percent of the children with autism, based on the brief observation alone.

"It's often not the pediatrician's fault that referrals are missed," Gabrielsen said. "Even autism experts missed a high percentage of referrals within that short timeframe. Decisions for referral need to be based on more information, including autism screening and information

from parents. We're hoping that this information can really empower parents to talk with pediatric care providers about their concerns."

The American Academy of Pediatrics recommends formal screening for autism at 18 and 24 months, but universal screening has not been fully adopted in pediatric primary care. The screening tools are also available for parents. These simple, standard tools for autism screening, such as the [M-CHAT-R checklist](#) and the CDC's [Learn the Signs. Act Early](#) campaign, are not perfect, but they are freely available, and can help parents learn what to look for as their child develops.

"Certainly, some young children with autism are clearly impaired and easy to recognize," said one of the other authors on the study Judith Miller. "However, this study looked at the entire range of children who present to the pediatrician's office, and we found that many children's impairments are not immediately obvious. For those children, formalized screening instruments and more time with a specialist may be critical."

Gabrielsen, an assistant professor in BYU's Counseling Psychology & Special Education department, and Miller, who is now at the Center for Autism Research at the Children's Hospital of Philadelphia, worked on the study with four other researchers. There were both school and clinical psychologists on the research team, and they worked in partnership with a large, independent community pediatrics clinic. The research began as a core group of autism experts at the University of Utah. The researchers are now affiliated with BYU; the Children's Hospital of Philadelphia; the University of Wisconsin-Madison's Waisman Center; Cleveland Clinic Children's; University of North Carolina, Chapel Hill, Asheville; and Tulane University.



Toddler Adelaide Burton is not too young to be screened for autism. A new study in Pediatrics shows that many her age aren't receiving referrals for further autism testing in pediatric exams.

Although there are still many unknowns about autism, one thing researchers do know is that early intervention makes a difference. If [autism](#) can be identified in the toddler years, intervention can begin while the brain is still rapidly developing and outcomes can change. A more comprehensive screening process, with parents and care providers working together, can have a great impact.

"Parents see their children at their very best and very worst," Gabrielsen said. "They're the experts for their children. They can be educated about

signs and symptoms, and need to help their care providers by speaking up if there's a problem and being involved in referral decisions."



Lynne Burton and daughter Adelaide Burton reading.

Provided by Brigham Young University

Citation: Brief observations miss autism symptoms in young children (2015, January 12)
retrieved 5 May 2024 from
<https://medicalxpress.com/news/2015-01-autism-symptoms-young-children.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.