

CRP/ESR disagreement common in infection, inflammation

January 16 2015



(HealthDay)—For patients with suspected infection or inflammation, C-reactive protein (CRP)/erythrocyte sedimentation rate (ESR) disagreement is common, according to a study published online Dec. 31 in the *International Journal of Rheumatic Diseases*.

Stephanie S bong, M.D., and Mark Feldman, M.D., from the Texas Health Presbyterian Hospital in Dallas, examined simultaneously ordered CRP and ESR results for 70 [adult patients](#) with suspected infection or [inflammation](#). The authors examined CRP/ESR disagreements and their stability in repeat testing in adults who had been tested on three separate occasions.

The researchers found that in 20 percent of [patients](#) there were initial

CRP/ESR disagreements. Of these 14 patients, only four had stable disagreements throughout the study, while 10 patients later exhibited CRP/ESR agreements. Of the 56 patients with initial CRP/ESR agreement, on subsequent testing, 10 developed CRP/ESR disagreements.

"Three mechanisms were identified to explain CRP/ESR disagreements: (i) slight fluctuations in the CRP and ESR around the upper limits of normal for these tests; (ii) development of an intercurrent illness; and (iii) different time courses of CRP and ESR elevations, in which the CRP rose and fell faster than the ESR," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: CRP/ESR disagreement common in infection, inflammation (2015, January 16)
retrieved 26 April 2024 from
<https://medicalxpress.com/news/2015-01-crpesr-disagreement-common-infection-inflammation.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--