

Earlier life adiposity trajectories linked to NAFLD in teens

January 8 2015



(HealthDay)—Earlier life trajectories of adiposity are associated with nonalcoholic fatty liver disease (NAFLD) in adolescents, according to a study published in the January issue of the *Journal of Gastroenterology and Hepatology*.

Oyekoya T. Ayonrinde, M.B.B.S., from the University of Western Australia in Perth, and colleagues examined the correlations between NAFLD diagnosed during adolescence and earlier life [trajectories](#) of anthropometry in a population-based cohort. Questionnaires and liver ultrasound were used to assess NAFLD in 1,170 adolescents aged 17 years.

The researchers found that 15.2 percent of adolescents were diagnosed with NAFLD. There was no association for birth anthropometry,

including birth weight, skinfold thickness, and ponderal index, with NAFLD. Adiposity differences between 17-year-olds with and without NAFLD were evident from age 3 years. There were correlations for greater adiposity trajectories for weight, [body mass index](#), skinfold thickness, mid-arm circumference, and chest circumference from age 3 onward with NAFLD diagnosis and severity of [hepatic steatosis](#) at age 17 years, particular in males (P

"Trajectories of childhood adiposity are associated with NAFLD," the authors write. "Exploration of clinically relevant risk factors and preventative measures for NAFLD should begin during childhood."

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: Earlier life adiposity trajectories linked to NAFLD in teens (2015, January 8) retrieved 4 May 2024 from
<https://medicalxpress.com/news/2015-01-earlier-life-adiposity-trajectories-linked.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
