

# High-dose statin may protect heart surgery patients' kidney health

January 20 2015

---

Acute kidney injury often arises after major surgery because the kidneys can be deprived of normal blood flow during the procedure. The use of contrast media, or dyes, can contribute to this problem.

In patients undergoing [coronary angiography](#) or [percutaneous coronary intervention](#), which are heart procedures that use dyes to help surgeons visualize the arteries, a high dose of the statin atorvastatin was linked with a reduction in blood levels of creatinine, a marker of kidney injury, as well as a lower incidence of [acute kidney injury](#) compared with a low dose of the statin.

The findings, which come from an analysis of 14 randomized controlled clinical trials consisting of 1689 patients, suggest that short-term high-dose atorvastatin therapy may be superior to low-dose atorvastatin for protecting heart surgery patients' kidney health. The findings are published in the *Journal of Clinical Pharmacology* study.

**More information:** Wu, H., Li, D., Fang, M., Han, H. and Wang, H. (2015), Meta-analysis of short-term high versus low doses of atorvastatin preventing contrast-induced acute kidney injury in patients undergoing coronary angiography/percutaneous coronary intervention. *Journal of Clinical Pharma*, 55: 123–131. [DOI: 10.1002/jcph.411](https://doi.org/10.1002/jcph.411)

Provided by Wiley

Citation: High-dose statin may protect heart surgery patients' kidney health (2015, January 20)  
retrieved 9 April 2024 from  
<https://medicalxpress.com/news/2015-01-high-dose-statin-heart-surgery-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.