


Infections increase death risk by 35 percent for ICU patients, study finds

January 5 2015



Effective infection prevention programs in the ICU can help elderly patients live longer and reduce the costs of care. Credit: Columbia University School of Nursing

Elderly patients admitted to intensive care units (ICUs) are about 35 percent more likely to die within five years of leaving the hospital if they develop an infection during their stay, a new study finds. Preventing two of the most common health care-acquired infections - sepsis caused by central lines and pneumonia caused by ventilators - can increase the odds that these patients survive and reduce the cost of their care by more than \$150,000.

"Any death from preventable infections is one too many," says senior study author Patricia Stone, PhD, RN, FAAN, director of the Center for Health Policy at Columbia University School of Nursing. "We've known for decades what works to prevent infections and save lives. Now, our study shows just how much money can be saved by investing in prevention."

The study, published in the January 2015 issue of the *American Journal of Infection Control*, looked at outcomes for 17,537 elderly Medicare patients admitted to 31 hospitals in 2002 to assess the cost and effectiveness of [infection prevention](#) efforts. Then, the researchers used an additional five years of Medicare claims data to assess the long-term outcomes and health costs attributed to health care-acquired infections.

While 57 percent of all the elderly ICU patients died within five years, the researchers found that infections made death more likely. For those who developed central line-associated bloodstream infections, or CLABSI, 75 percent died within five years, as did 77 percent of those who developed ventilator-associated pneumonia, or VAP.

Effective prevention programs for CLABSI resulted in an estimated gain of 15.55 years of life on average for all patients treated in the ICU, the study found. Catheters, also known as a central line, deliver life-saving medicines and nutrition. Without proper insertion, utilization, and maintenance, catheters can also transmit deadly infections to the bloodstream. Simple [infection](#) prevention measures include hand washing before handling the catheter and immediately changing the dressing around the central line if it gets wet or dirty.

Efforts to prevent VAP resulted in an estimated gain of 10.84 years of life on average for all patients treated in the ICU. Ventilators, machines that force air into the lungs when patients can't breathe on their own, can cause infections when patients lie flat in bed for long periods of time.

Keeping [patients](#) elevated in bed, with the head higher than the feet, is one simple precaution that can help prevent pneumonia.

On average, the ongoing cost of running an infection prevention program in the ICU is about \$145,000, the study found. Prevention efforts reduced ICU costs by \$174,713 per patient for each instance of CLABSI, and by \$163,090 for VAP.

"This evidence points definitively to the value of investing in infection prevention," says lead study author Andrew Dick, PhD, senior economist at RAND Corporation.

Health care-associated infections kill an estimated 100,000 Americans a year and create approximately \$33 billion in excess medical costs. The U.S. Centers for Disease Control and Prevention first linked infection rates to prevention programs in the 1970s. Research since then has shown that checklists and other targeted [infection control](#) practices can make a significant dent in [infection rates](#).

More information: The paper is titled "A decade of investment in infection prevention: A cost-effectiveness analysis."

Provided by Columbia University Medical Center

Citation: Infections increase death risk by 35 percent for ICU patients, study finds (2015, January 5) retrieved 23 April 2024 from <https://medicalxpress.com/news/2015-01-infections-death-percent-icu-patients.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--