

# Interferon-free combination therapy prevents recurrence of HCV after liver transplantation

January 9 2015

---

A 24-week course of sofosbuvir and ribavirin can eliminate hepatitis C virus (HCV) infection in two-thirds of patients who undergo liver transplantation, with positive consequences on their short- and long-term prognoses, according to two new studies published in *Gastroenterology*, the official journal of the American Gastroenterological Association.

The [first study](#), a Phase 2, open-label study, found that combination therapy with sofosbuvir and ribavirin for up to 48 weeks before liver transplantation can prevent recurrence of HCV infection post-transplantation in 70 percent of patients. This study provides proof of concept that successful interferon-free treatment prior to liver removal can eliminate the disease in most patients.

"Patients with hepatitis C virus at the time of liver transplantation universally experience recurrent HCV infection," said lead study author Michael P. Curry, MD, from Beth Israel Deaconess Medical Center, Boston, MA. "Recurrent HCV infection follows an aggressive course. Given the burden of disease—the increased morbidity, mortality and costs—and the lack of a safe and broadly effective treatment to prevent recurrence of HCV infection, these results provide hope for patients in need."

The [second study](#) was a prospective, multi-center, open-label pilot study, consisting of patients with recurrent HCV infection after a primary or secondary [liver transplant](#). Patients received an all-oral regimen of sofosbuvir and ribavirin for 24 weeks, which resulted in sustained

virologic response, with no detectable virus, in 70 percent of patients. This rate is similar to that achieved in clinical trials using the same dose and duration of sofosbuvir and ribavirin in the nontransplant setting.

"A well tolerated and effective treatment protocol for recurrence of HCV [infection](#) following [liver transplantation](#) is an important unmet clinical need," said lead study author Michael R. Charlton, MD, Mayo Foundation, Rochester, MN. "Our study demonstrates that [patients](#) with characteristics that have historically been difficult to cure with interferon-based regimens, including those with advanced disease, may benefit from this all-oral interferon-free therapy."

In both studies, the rate of discontinuation due to adverse events was low.

**More information:** Curry, M.P., et al. Sofosbuvir and Ribavirin Prevent Recurrence of HCV Infection after Liver Transplantation: An Open-Label Study, *Gastroenterology*, Volume 148(1): 100-107.e1

Charlton, M.R., et al., Sofosbuvir and Ribavirin for Treatment of Recurrent Hepatitis C Virus Infection After Liver Transplantation, *Gastroenterology*, Volume 148(1): 108-117

Provided by American Gastroenterological Association

Citation: Interferon-free combination therapy prevents recurrence of HCV after liver transplantation (2015, January 9) retrieved 23 April 2024 from <https://medicalxpress.com/news/2015-01-interferon-free-combination-therapy-recurrence-hcv.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.