

# Study of former NFL players reveals specifics of concussive brain damage

January 26 2015

---

A team of Johns Hopkins specialists, using a battery of imaging and cognitive tests, has gathered evidence of accumulated brain damage that could be linked to specific memory deficits in former National Football League (NFL) players experienced decades after they stopped playing the game.

Results of the small study of nine men provide further evidence for potential long-term neurological risk to football [players](#) who sustain repeated concussions and support calls for better player protections.

"We're hoping that our findings are going to further inform the game," says Jennifer Coughlin, M.D., assistant professor of psychiatry and behavioral sciences at the Johns Hopkins University School of Medicine. "That may mean individuals are able to make more educated decisions about whether they're susceptible to [brain injury](#), advise how helmets are structured or inform guidelines for the game to better protect players."

Several anecdotal accounts and studies have suggested that athletes, such as collegiate and professional football, hockey, and soccer players, exposed to repeat concussions could suffer permanent [brain damage](#) and deficits from these events. However, the mechanism of damage and the source of these deficits have been unclear.

To reveal them, Coughlin; Yuchuan Wang, Ph.D., assistant professor of radiology and radiological science at the Johns Hopkins University School of Medicine; and their colleagues used tests to directly detect

deficits and to quantify localized molecular differences between the brains of former players and healthy people who didn't play football.

The researchers recruited nine former NFL players who retired decades ago and ranged in age from 57 to 74. The men had played a variety of team positions and had a wide range of self-reported, historical concussions, varying from none for a running back to 40 for a defensive tackle. The researchers also recruited nine age-matched "controls"—healthy individuals who had no reason to suspect they had brain injuries.

Each of the volunteers underwent a positron emission tomography (PET) scan, a test in which an injected radioactive chemical binds to a specific biological molecule, allowing researchers to physically see and measure its presence throughout the body. In this case, the research team focused on the translocator protein, which signals the degree of damage and repair in the brain. While healthy individuals have low levels of this protein spread throughout the brain, those with brain injuries tend to have concentrated zones with high levels of translocator protein wherever an injury has occurred.

The volunteers also underwent MRIs, which allowed the researchers to match up the PET scan findings with anatomical locations in the volunteers' brains and check for structural abnormalities. In addition, they took a battery of memory tests.

While the control volunteers' tests showed no evidence of brain damage, PET scans showed that on average, the group of former NFL players had evidence of brain injury in several temporal medial lobe regions, including the amygdala, a region that plays a significant role in regulating mood. Imaging also identified injuries in many players' supramarginal gyrus, an area linked to verbal memory.

While the hippocampus, an area that plays a role in several aspects of memory, didn't show evidence of damage in the PET scans, MRIs of the former players' brains showed atrophy of the right-side hippocampus, suggesting that this region may have shrunk in size due to previous damage.

Additionally, many of the NFL players scored low on memory testing, particularly in tests of verbal learning and memory.

Though the researchers emphasize that this pilot study, published in the February 2015 issue of the journal *Neurobiology of Disease*, is small in size, they say that the evidence among just nine former NFL players suggests that there are molecular and structural changes in specific brain regions of athletes who have a history of repetitive hits to the head, even many years after they've left active play.

The researchers are currently looking for translocator protein hotspots in both active and recently retired players to help determine whether these changes develop close to the time of play or whether they're a result of a more delayed response to injury with similarities to other degenerative brain disorders.

If these findings are seen in studies with larger numbers of participants, they say, use of this molecular [brain](#) imaging technique could eventually lead to changes in the way players are treated after experiencing concussion or how contact sports are played.

Provided by Johns Hopkins University School of Medicine

Citation: Study of former NFL players reveals specifics of concussive brain damage (2015, January 26) retrieved 19 April 2024 from <https://medicalxpress.com/news/2015-01-nfl-players-reveals-specifics-concussive.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.