

# Obesity and diabetes symptoms in mice improved by reversing brain inflammation

January 29 2015

---

Using an antioxidant to reverse inflammation in the brain caused by a high-fat diet greatly improves symptoms related to obesity and type II diabetes, a new study from New Zealand's University of Otago suggests.

The research, which appears in the leading international journal *Diabetes*, was led by Dr Alex Tups of the University's Centre for Neuroendocrinology and Department of Physiology.

Dr Tups and an international team investigated whether directly stopping [inflammatory processes](#) in the brain's hypothalamus could help lower [blood sugar levels](#) and reduce [insulin resistance](#).

In their research the team blocked a particular inflammatory signalling pathway (IKK $\beta$ /NF- $\kappa$ B) in the brains of obese [mice](#). The researchers studied both mice that were obese due to a deficiency in the satiety hormone leptin and others due to a high-fat diet.

The scientists administered butein to the mice to block the signalling pathway, which is involved in the body's inflammatory immune responses. Butein is a flavonoid derived from plants traditionally used in Chinese herbal medicine.

Dr Tups says the team found that administering butein either directly into the brain or orally greatly improved glucose tolerance and brain insulin signalling in both types of [obese mice](#).

"We also showed that this profound effect was dose-dependent with better glucose tolerance achieved through higher doses of butein," Dr Tups says.

The improved [glucose tolerance](#) of high-fat diet mice treated with the antioxidant was such that no difference was noticeable between them and low fat-diet mice that had not received butein.

To confirm that activation of the IKK $\beta$ /NF- $\kappa$ B pathway plays a central role in metabolic obesity symptoms, the researchers also used a gene therapy technique to inhibit it in neurons in the hypothalamus.

This gene therapy resulted in high-fat diet mice having a reduced body weight, building up less fat, expending more energy, and showing evidence of improved leptin-signalling.

Dr Tups says the study adds to growing body of evidence that a diet high in saturated fats activates a cascade of inflammatory processes in the brain which impair leptin and insulin signalling, leading to obesity and type II diabetes.

"Our findings strongly support this idea and we also show that reversing this inflammation promotes a return towards normal metabolic functioning," he says.

The research suggests that butein and other natural compounds that block inflammation in the brain should be vigorously investigated as novel anti-diabetic treatments, he says.

Provided by University of Otago

Citation: Obesity and diabetes symptoms in mice improved by reversing brain inflammation

(2015, January 29) retrieved 2 May 2024 from <https://medicalxpress.com/news/2015-01-obesity-diabetes-symptoms-mice-reversing.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.