


# Prescription painkillers, widely used by childbearing age women, double birth defects risk

January 22 2015



Steps women can take to help give her baby a healthy start in life. Credit: March of Dimes

More than one-fourth of privately-insured and one-third of Medicaid-enrolled women of childbearing age filled prescriptions for opioid-based (narcotic) painkillers between 2008 and 2012, according to a new analysis published today by the U.S. Centers for Disease Control and Prevention (CDC) in its *Morbidity and Mortality Weekly Report* (MMWR).

Many women are unaware that prescription opioid-based medications such as codeine, oxycodone, hydrocodone, or morphine, used to treat severe pain, may increase the risk for serious birth defects of the baby's brain, spine, and heart, as well as preterm birth when taken during pregnancy. Use of these medications also can cause babies to suffer withdrawal symptoms when born, a condition known as [neonatal abstinence syndrome](#) or NAS, a growing problem in U.S. birthing hospitals.

Since half of all pregnancies are unplanned, women may be prescribed opioid-based pain medications before they or their [health care providers](#) know they are pregnant. "This highlights the importance of promoting safer alternative treatments, when available for women of reproductive age. We must do what we can to protect babies from exposure to opioids." stated Coleen A. Boyle, PhD, MSHyg, Director of CDC's National Center on Birth Defects and Developmental Disabilities (NCBDDD). "

"If you are using an opioid pain killer, you should also be practicing effective birth control," says José F. Cordero, MD, MPH, a pediatrician,

birth defects expert formerly at CDC, and member of the March of Dimes Board of Trustees. "If you decide to get pregnant or do become pregnant, tell your [health care](#) provider about all the medications you are taking right away. You may be able to switch to a safer alternative." Dr. Cordero also urged physicians and other prescribers not to write prescriptions for opioid-based painkillers for their female patients who may become pregnant without a discussion of the risks and safer alternatives.

January is Birth Defects Prevention Month and the CDC and the March of Dimes are taking this opportunity to raise awareness about the risk of birth defects from opioid-based prescription pain medications during pregnancy.

In the U.S., a baby is born with a birth defect every four and a half minutes, and one out of every five deaths in the first year of life is caused by a [birth defect](#). In addition to the human toll, birth defects incur hospital-related economic costs that exceed \$2.6 billion annually.

"The CDC's Treating for Two: Safer Medication Use in Pregnancy initiative offers information to women and their healthcare providers about medication use during pregnancy. This initiative aims to prevent birth defects and improve the health of mothers by working to identify the best alternatives for treatment of common conditions during pregnancy and during the childbearing years," explains Dr. Boyle.

**More information:** [www.cdc.gov/treatingfortwo](http://www.cdc.gov/treatingfortwo)

Provided by March of Dimes Foundation

Citation: Prescription painkillers, widely used by childbearing age women, double birth defects

risk (2015, January 22) retrieved 3 May 2024 from  
<https://medicalxpress.com/news/2015-01-prescription-painkillers-widely-childbearing-age.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.