


Rare amino acid influences E. coli infection

January 12 2015



Escherichia coli. Credit: Rocky Mountain Laboratories, NIAID, NIH

Scientists have discovered how a rare amino-acid in humans influences the behavior of the E.coli bacterium.

Most of the thousands of strains of E. coli are harmless, with many being

a normal part of the [gut flora](#) in [healthy people](#), however some strains can cause illness in humans.

Among the most well-known is E. coli O157, typically acquired via contaminated food, which causes severe diarrhea and can lead to [kidney damage](#).

The O157 strain only infects the gut so scientists at the University of Glasgow wanted to know what stopped it from spreading to other parts of the body.

The team led by Dr Andrew Roe, and PhD student James Connolly of the Institute of Infection, Immunity & Inflammation, analyzed the genome sequence of 1,500 strains of E.coli.

They wanted to see how the genes of the bug, which enable it to attach to and infect a host, responded to varying concentrations of D-Serine, an amino-acid produced in the brain where it plays a role in nerve signaling.

They found that E. coli O157 is unable to attach itself to host tissue in high concentrations of D-Serine. Other strains, such as those that cause meningitis, thrive in the presence of the amino-acid.

The discovery, published in the *ISME Journal*, opens up the possibility of altering the diet to increase levels of D-Serine to prevent E. coli O157 infection or perhaps treat it.

Dr Andrew Roe, senior lecturer, said: "This work provides new insights into the infection process with the aim of developing compounds that block such bugs from attaching to the host.

"With many strains of E. coli developing resistance to traditional antibiotics, such approaches are urgently needed.

"If we can disarm such bacteria rather than killing them it puts less pressure on the bacteria to evolve into something that is resistant to treatment."

E. coli O157 doesn't normally live in humans, instead residing in the gut of cattle. Eating contaminated food is the most common cause of infection but it can also be picked up in the environment, through contact with the bacteria in fields, for example.

The genetic variety between strains of *E. coli* is huge, with around 2,000 'core' genes and 18,000 genes that vary between strains. Different [strains](#) are able to attach themselves to different tissues, causing a range of different infections.

The bacterium can cause a wide range of infections including those of the gut, bladder, bloodstream and brain. These can be very common, for example, over half of all women suffer from *E. coli* associated bladder infections at some point in their lives.

More information: Link to research paper in The *ISME Journal*:
[www.nature.com/ismej/journal/v ... /ismej2014242a.html](http://www.nature.com/ismej/journal/v.../ismej2014242a.html)

Provided by University of Glasgow

Citation: Rare amino acid influences *E. coli* infection (2015, January 12) retrieved 25 April 2024 from <https://medicalxpress.com/news/2015-01-rare-amino-acid-coli-infection.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.