


Scientists call for antibody 'bar code' system to follow Human Genome Project (Update)

February 4 2015



Structure of the human antibody molecule with heavy chain in blue and light chain in red (IgG isotype shown). Credit: Garvan Institute of Medical Research / Centre for Targeted Therapy

More than 100 researchers from around the world have collaborated to craft a request that could fundamentally alter how the antibodies used in research are identified, a project potentially on the scale of the now-completed Human Genome Project.

"We propose that antibodies be defined by their sequences, just as genes are," said Andrew Bradbury, a researcher at Los Alamos National

Laboratory, "and they should be made recombinantly in cell lines."

Referring to antibodies according to the sequences encoding their various subunits, their concentrations and the standardized experimental buffers used for each assay would enable researchers world-wide to employ the same affinity reagents under the same conditions.

The sequence of an antibody or binding reagent, is the ultimate "bar code" for that reagent, ensuring that everyone can use the same reagent for the same target. Deriving the bar code involves either selecting antibodies from in vitro libraries, or cloning and sequencing antibody genes from hybridomas, the cells that traditionally make monoclonal antibodies. However, it will require a paradigm shift in the way antibodies are supplied.

Antibodies are the specialized proteins that normally help the body identify and neutralize bacterial and other attacks on the immune system, and have been exploited by scientists as specific binding reagents. The problem that the researchers seek to resolve with their new sequencing and recombinant expression demand is that of antibody quality control and accurate identification. Antibodies are the only widely used reagents in biological research that are not defined at the sequence level (unlike genes, oligos, plasmids, recombinant proteins, etc.), Bradbury noted.

The quality of all antibodies produced varies enormously according to the manufacturer, and most antibodies are rarely validated and between-batch differences are common, the researchers note.

Also, the quality of the documentation accompanying the batches (for instance, on the concentration of antibody present in each vial) is enormously variable; even when it is provided, it often doesn't correspond to the batch being supplied, the letter says.

"To stem the enormous waste in materials, researcher time and money caused by their lack of validation and characterization, antibodies must be defined by their sequences, and in the main, produced de novo under standardized conditions," the writers point out. "De novo" production in this case would refer to the process of generating antibodies or other binding reagents from scratch using methods that do not involve animal immunization, rather than cloning antibodies from cells derived from immunization.

For polyclonal antibodies, for example, production is based on injecting an animal with a protein and then later extracting the antibodies that the animal's blood is carrying in response to the immunizing protein. The varied antibodies are anything but precise, however, with only 0.5 to 5 percent of the product being the desired antibodies, ready to respond to the original target. The rest are antibodies already in the animals' blood stream, and represent previous immunological challenges. *Nature* is simply not precise in this process, and the scattershot harvesting of so many of the wrong antibodies is producing waste in the research pipeline, by producing antibodies with poor specificity.

"What we are proposing is to move over to an in vitro method of generating antibodies that will not require the use of animals at all, and importantly, will directly lead to molecules with known sequences," Bradbury said.

The scores of researchers signing the letter are united in their desire for a more structured, accurate antibody identification system, although Bradbury and his cohorts recognize that significant costs will be associated with the effort.

"The National Institutes of Health and European Union were very proactive in starting pilot programs to address part of this issue, and given the long-term investment required, we wanted to encourage them

to continue," he said. "Given advances in the generation of specific binding reagents by in vitro methods, and the recent discussions on reproducibility in science, we thought it was about time this issue was addressed."

To make uniform reagents in the laboratory for all 20,000 human gene products, the authors estimate that using current technology, an investment of roughly a billion dollars would be required. "This is less than the money wasted worldwide on bad reagents in under two years," Bradbury said, and it would be recovered in time through the production of better research results, the team suggests. However, the total amount required would be expected to be far less if appropriate investments in technology development were made.

More information: *Nature* Comment, [DOI: 10.1038/518027a](https://doi.org/10.1038/518027a)"
target="_blank">nature.com/articles/[DOI: 10.1038/518027a](https://doi.org/10.1038/518027a)

Provided by University of Zurich

Citation: Scientists call for antibody 'bar code' system to follow Human Genome Project (Update) (2015, February 4) retrieved 26 April 2024 from <https://medicalxpress.com/news/2015-02-antibodies-lab-higher-quality-dna.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
