

# Certain macrolides linked with higher risk of pyloric stenosis

February 17 2015

---



(HealthDay)—New research supports previous findings that erythromycin can increase the risk of infantile hypertrophic pyloric stenosis (IHPS). The research also indicates that azithromycin is associated with a higher risk of IHPS when given to infants under 6 weeks old. The findings were published online Feb. 16 in the *Pediatrics*.

Matthew D. Eberly, M.D., from the Uniformed Services University of the Health Sciences in Bethesda, Md., and colleagues analyzed the records of 1,074,236 children and compared those who received [erythromycin](#) or [azithromycin](#) in their first three months to those who didn't. Overall, 2,466 [infants](#) developed IHPS and 6,777 infants had received one of the two antibiotics studied. The overall risk of IHPS was small—occurring in 2.29 babies per 1,000.

The researchers found that the use of erythromycin was associated with a 13.3-fold increased risk of IHPS in the first two weeks of life, and a 4.1-fold higher risk when used between the ages of 2 and 6 weeks. The team also found that azithromycin was associated with an 8.26-fold higher risk when used during the first two weeks of life, and a 2.98-fold higher risk when used in infants between the ages of 2 and 6 weeks.

"Ingestion of oral azithromycin and erythromycin places young infants at increased risk of developing IHPS," the authors write. "This association is strongest if the exposure occurred in the first two weeks of life, but persists to a lesser degree in children between 3 and 6 weeks of age."

**More information:** [Abstract](#)  
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: Certain macrolides linked with higher risk of pyloric stenosis (2015, February 17)  
retrieved 2 May 2024 from  
<https://medicalxpress.com/news/2015-02-macrolides-linked-higher-pyloric-stenosis.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--