


New method for minimally invasive tissue ablation surgery

February 6 2015



The increase in the extent of tissue ablation achieved by combining reversible electroporation pulses with electrolysis is shown with Masson's trichromatic staining of the liver tissue. Plate electrodes were placed on the top and bottom surfaces of the rat liver lobe in vivo. An electrolytic protocol consisting of 4 mA of current applied over a 30s time interval produced tissue ablation around the top and bottom surfaces of the tissue that were in contact with the electrodes, as shown in panel (a). A reversible electroporation protocol (applying an electric field of 250 V/cm over 8 pulses of 100 μ s in length at a 1 Hz frequency) also resulted in tissue ablation concentrated near the top and bottom surfaces of the

tissue (b). Combining electrolysis with reversible electroporation, however, resulted in a much greater region of tissue ablation. The treatment zone is seen to penetrate through the entire liver thickness, as outlined in panels (c) and (d). In panel (c), the electrolysis protocol was delivered first, followed by the reversible electroporation protocol, and in panel (d), the reversible electroporation protocol was applied before electrolysis. Dashed lines are used to highlight the treated area. Credit: *TECHNOLOGY* journal.

A team of researchers that includes scientists from the Quinnipiac University and the University of California, Berkeley reports a new method for minimally invasive tissue ablation surgery that combines electrolysis with reversible electroporation.

The interaction between electrical currents and biological media produces a variety of different biophysical phenomena, including two relevant to this study, electroporation and [electrolysis](#). Electroporation occurs when very brief, microsecond length, high-magnitude electric fields are applied across cells, causing pores to form in the cell membrane. In reversible electroporation these pores are transient, and most cells survive the process. Since the early 1980's work of Neumann, reversible electroporation has become a routine method for introducing foreign genes and other molecules into cells. Tissue ablation by electrolysis emerged from the early 19th century work of Faraday on the chemical species produced at electrode surfaces immersed in an ionic solution. These chemical species diffuse away from the electrodes and into the tissue, causing local changes in pH. The resulting acidic and basic regions, as well as the presence of some of these new chemical species, causes cell death. This mechanism of cell death has been employed for [minimally invasive surgery](#) since the early 1800's. Electrolytic ablation is seen as advantageous since it requires very low currents and voltages. However, it is a time consuming process, because the method of cell death involves rate and dose dependent chemical

reactions. The length of the procedure is considered a major disadvantage of electrolytic ablation, which has made it less attractive to surgeons.

By applying electric currents that modulate electrolysis with reversible electroporation pulses, the research group has shown that the tissue ablation by electrolysis is greatly increased, resulting in a new, more effective tissue ablation technique. The first study of this new ablation method is set to appear in the upcoming issue of the journal *TECHNOLOGY*.

The team hypothesizes that adding reversible electroporation pulses makes electrolysis so much more effective, because the pores formed in the cell membranes by reversible electroporation allow the [chemical species](#) produced by electrolysis to enter into the cell, causing [cell death](#) at much lower concentrations. By adding a sequence of reversible electroporation pulses, electrolysis may be able to be used to ablate larger cancer tumors. In addition, this new method may provide greater control over the region of tissue that gets treated, resulting in safer and more effective cancer treatment.

Provided by World Scientific Publishing

Citation: New method for minimally invasive tissue ablation surgery (2015, February 6) retrieved 3 May 2024 from

<https://medicalxpress.com/news/2015-02-method-minimally-invasive-tissue-ablation.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--