

Napping reverses health effects of poor sleep

February 10 2015

A short nap can help relieve stress and bolster the immune systems of men who slept only two hours the previous night, according to a new study published in the Endocrine Society's *Journal of Clinical Endocrinology & Metabolism (JCEM)*.

Lack of [sleep](#) is recognized as a public health problem. Insufficient sleep can contribute to reduced productivity as well as vehicle and industrial accidents, according to the U.S. Centers for Disease Control and Prevention. In addition, people who sleep too little are more likely to develop chronic diseases such as obesity, diabetes, high blood pressure and depression.

Nearly three in 10 adults reported they slept an average of six hours or less a [night](#), according to the National Health Interview Survey.

"Our data suggests a 30-minute nap can reverse the hormonal impact of a night of poor sleep," said one of the *JCEM* study's authors, Brice Faraut, PhD, of the Université Paris Descartes-Sorbonne Paris Cité in Paris, France. "This is the first study that found napping could restore biomarkers of neuroendocrine and immune health to normal levels."

The researchers used a cross-over, randomized study design to examine the relationship between hormones and sleep in a group of 11 healthy men between the ages of 25 and 32. The men underwent two sessions of sleep testing in a laboratory, where meals and lighting were strictly controlled.

During one session, the men were limited to two hours of sleep for one night. For the other session, subjects were able to take two, 30-minute naps the day after their sleep was restricted to two hours. Each of the three-day sessions began with a night where subjects spent eight hours in bed and concluded with a recovery night of unlimited sleep.

Researchers analyzed the participants' urine and saliva to determine how restricted sleep and napping altered hormone levels. After a night of limited sleep, the men had a 2.5-fold increase in levels of norepinephrine, a hormone and neurotransmitter involved in the body's fight-or-flight response to [stress](#). Norepinephrine increases the body's heart rate, blood pressure and blood sugar. Researchers found no change in norepinephrine levels when the [men](#) had napped following a night of limited sleep.

Lack of sleep also affected the levels of interleukin-6, a protein with antiviral properties, found in the subjects' saliva. The levels dropped after a night of restricted sleep, but remained normal when the subjects were allowed to nap. The changes suggest naps can be beneficial for the [immune system](#).

"Napping may offer a way to counter the damaging effects of sleep restriction by helping the immune and neuroendocrine systems to recover," Faraut said. "The findings support the development of practical strategies for addressing chronically sleep-deprived populations, such as night and shift workers."

Other authors of the study include: Samir Nakib, Catherine Drogou, Maxime Elbaz, Fabien Sauvet, Jean-Pascal De Bandt and Damien Léger of the Université Paris Descartes-Sorbonne Paris Cité.

More information: "Napping Reverses the Salivary Interleukin-6 and Urinary Norepinephrine Changes Induced by Sleep Restriction," *Journal*

of Clinical Endocrinology & Metabolism, 2015.

Provided by The Endocrine Society

Citation: Napping reverses health effects of poor sleep (2015, February 10) retrieved 26 April 2024 from <https://medicalxpress.com/news/2015-02-napping-reverses-health-effects-poor.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.