

Review: Some nonpharmacologic Tx effective in peds GI disorders

February 9 2015



(HealthDay)—Certain nonpharmacologic treatments are effective in pediatric abdominal pain-related functional gastrointestinal disorders (AP-FGIDs), according to a review published online Feb. 9 in *Pediatrics*.

Juliette M.T.M. Rutten, M.D., from Emma's Children's Hospital in Amsterdam, and colleagues conducted a systematic review of the literature relating to nonpharmacologic interventions for pediatric AP-FGIDs. Data were included from 24 randomized controlled trials that included 1,390 children.

The researchers observed significant improvement of [abdominal pain](#) after hypnotherapy versus standard care/wait list approaches and after [cognitive behavioral therapy](#) versus control/wait list approaches. At the

six-month follow-up only, written self-disclosure improved pain frequency. Significantly more treatment responders were seen with *Lactobacillus rhamnosus* GG (LGG) and VSL#3 versus placebo (LGG relative risk, 1.31; VSL#3 P

"Although high-quality studies are lacking, some evidence shows efficacy of hypnotherapy, cognitive behavioral therapy, and probiotics (LGG and VSL#3) in pediatric AP-FGIDs," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: Review: Some nonpharmacologic Tx effective in peds GI disorders (2015, February 9)
retrieved 6 May 2024 from
<https://medicalxpress.com/news/2015-02-nonpharmacologic-tx-effective-peds-gi.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
