

Proton pump inhibitor use not linked to cardiac arrhythmia

February 24 2015



In critically ill patients, proton pump inhibitor use is not associated with cardiac arrhythmia, according to a study published online Feb. 6 in the *Journal of Clinical Pharmacology*.

(HealthDay)—In critically ill patients, proton pump inhibitor (PPI) use is not associated with cardiac arrhythmia, according to a study published online Feb. 6 in the *Journal of Clinical Pharmacology*.

Kenneth P. Chen, M.D., from the Beth Israel Deaconess Medical Center in Boston, and colleagues examined whether PPI exposure correlated with admission electrocardiogram readings of a [cardiac arrhythmia](#) in a study involving more than 8,000 [critically ill patients](#).

Within the cohort there were 25.4 percent PPI users and 6 percent taking a histamine 2 antagonist. Of the patients, 14.0 percent had a cardiac arrhythmia. The researchers found that the risk of arrhythmia in

association with PPI use was 1.15 (95 percent confidence interval, 1.00 to 1.32; $P = 0.04$) in unadjusted analyses and 0.91 (95 percent confidence interval, 0.77 to 1.06; $P = 0.22$) in adjusted analyses. PPI use was not associated with increased risk of cardiac arrhythmia among diuretic users (2,476 patients).

"In summary, in a large cohort of critically ill patients, PPI exposure is not associated with an increased risk of cardiac arrhythmia," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: Proton pump inhibitor use not linked to cardiac arrhythmia (2015, February 24)
retrieved 26 April 2024 from
<https://medicalxpress.com/news/2015-02-proton-inhibitor-linked-cardiac-arrhythmia.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--