

# Ranibizumab reverses vision loss caused by diabetes

February 18 2015

---

Ranibizumab, a prescription drug commonly used to treat age-related vision loss, also reverses vision loss caused by diabetes among Hispanic and non-Hispanic whites, according to a new study led by investigators from the University of Southern California (USC) Eye Institute.

Diabetic retinopathy and [diabetic macular edema](#) are the leading causes of [vision loss](#) in working-age adults in the United States, according to the National Eye Institute. Laser surgery is the standard treatment for advanced stages of the disease, characterized by blurred vision, but previous research has shown that only 30 percent of patients saw improvement in their vision.

"We found that ranibizumab can save the sight of thousands of working-age individuals suffering from [diabetic eye disease](#), as standard treatments such as laser are not as effective," said Rohit Varma, M.D., M.P.H., director of the USC Eye Institute, professor and chair of ophthalmology at the Keck School of Medicine of USC and the study's lead author.

Varma's team developed a population-based model that suggests that administering 0.3 milligrams of ranibizumab every four weeks to patients with diabetic macular edema would reduce the number of cases of vision impairment by 45 percent, or 5,134 individuals, and the number of cases of legal blindness by 75 percent, or 1,275 individuals. The model was based on the approximately 37,000 Hispanic and non-Hispanic white adults with diabetic macular edema in the United States

for whom ranibizumab treatment could be used. Because other race and ethnic groups were not included in the study, authors contend that the treatment may benefit even more people than their results show.

Ranibizumab is manufactured and marketed by Genentech Inc. under the trade name Lucentis. The study was supported in part by Genentech Inc. It appears in the Feb. 7, 2015, online edition of the peer-reviewed medical journal *Ophthalmology*.

**More information:** Varma, R., Bressler, N. M., Doan, Q. V., Danese, M., Dolan, C. M. ... Turpcu, A. (2015). Visual impairment and blindness avoided with ranibizumab in Hispanic and non-Hispanic whites with diabetic macular edema in the United States. *Ophthalmology*, 1-8. Published online Feb. 9, 2015; doi:10.1016.j.optha.2014.007

Provided by University of Southern California

Citation: Ranibizumab reverses vision loss caused by diabetes (2015, February 18) retrieved 19 April 2024 from <https://medicalxpress.com/news/2015-02-ranibizumab-reverses-vision-loss-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.