

# Salmonella and Campylobacter show significant levels of resistance to common antimicrobials in humans and animals

February 28 2015

---

Treatment options for some of the most common food-borne infections are decreasing, as types of bacteria (called 'isolates') continue to show resistance to antimicrobial drugs. For example, multi-drug resistant isolates of *Salmonella* continue to spread across Europe. Also, high resistance to the antimicrobial ciprofloxacin in *Campylobacter* isolates in both humans and animals has been reported in some Member States. Encouragingly, co-resistance to critically important antimicrobials for both bacteria remains low. These are some the findings of the latest EFSA-ECDC European Union Summary Report on antimicrobial resistance (AMR) in zoonotic and indicator bacteria from humans, animals and food, which analyses data from 2013.

For the first time, EFSA and ECDC have used similar criteria to interpret data. "Findings in [antimicrobial resistance](#) in humans, animals and foods are now more comparable. This is a step forward in the fight against antimicrobial resistance", said Marta Hugas, Acting Head of EFSA's Risk Assessment and Scientific Assistance Department.

"The high levels of resistance to fluoroquinolones observed in *Campylobacter* isolates from both humans and broilers are of concern considering that a large proportion of human *Campylobacter* infections come from handling, preparation and consumption of broiler meat. Such high resistance levels reduce the effective treatment options for severe human *Campylobacter* infections", said Mike Catchpole, Chief Scientist

at ECDC.

## Key findings

- Resistance in *Salmonella* to commonly used antimicrobials was frequently detected in humans and animals (especially broilers and turkeys) and derived meat products. Multi-drug resistance was high (in humans 31.8%, in broilers 56.0%, in turkey 73.0%, and in fattening pigs 37.9%), and the continued spread of particularly multi-drug resistant clones reported in both human and animal (broilers, pigs and cattle) isolates is of concern.
- Resistance to commonly used antimicrobials in *Campylobacter* isolates was frequently detected in humans and animals (especially broilers, pigs and cattle). In food, resistance was detected in broiler meat. Resistance to ciprofloxacin, a critically important antimicrobial, was particularly high in humans (meaning that treatment options for serious infections with these zoonotic bacteria are reduced). In *Campylobacter jejuni* more than half of both human and broiler isolates (54.6% and 54.5% respectively) were resistant, alongside 35.8% in cattle. In *C. coli* two thirds of humans and broiler isolates (66.6% and 68.8% respectively) were resistant along with 31.1% of pig isolates.
- Levels of co-resistance to critically important antimicrobials in *Salmonella* were low (in humans 0.2%, in broilers 0.3%, and in fattening pigs and in turkey there was none). Levels of multi-drug resistance and co-resistance in *Campylobacter* isolates to critically important antimicrobials were generally reported at low to moderate levels in animals (in *C. jejuni* isolates from broilers and cattle 0.5% and 1.1%, respectively, in *C. coli* isolates from broilers and fattening pigs 12.3% and 19.5%, respectively) and at low levels in humans (1.7% in *C. jejuni* and 4.1% in *C. coli*).

The report also includes data on resistance in indicator *Escherichia coli*,

*indicator enterococci* and methicillin-resistant *Staphylococcus aureus*, in animals and food.

**More information:** [www.efsa.europa.eu/en/efsajournal/pub/4036.htm](http://www.efsa.europa.eu/en/efsajournal/pub/4036.htm)

Provided by European Food Safety Authority

Citation: Salmonella and Campylobacter show significant levels of resistance to common antimicrobials in humans and animals (2015, February 28) retrieved 3 May 2024 from <https://medicalxpress.com/news/2015-02-salmonella-campylobacter-significant-resistance-common.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.