

Simple test detects increased risks in patients with acute kidney injury

February 5 2015

A simple test can help predict which patients with acute kidney injury will likely develop serious kidney problems and die in the hospital, according to a study appearing in an upcoming issue of the *Journal of the American Society of Nephrology* (JASN). Using this test could help clinicians safeguard patients' health.

Acute kidney injury (AKI) is an abrupt decline in kidney function that often arises after major surgeries or severe infections. When a patient develops AKI, some may recover promptly while others get worse and require dialysis. Unfortunately, there are limited tools to help predict which patients with early AKI will go on to develop the more severe forms of AKI that require dialysis. Having such information would improve care and perhaps help safeguard patients' health.

In a study of 77 patients with early AKI, a team led by Lakhmir Chawla, MD (George Washington University and Veteran Affairs Medical Center, Washington DC) and Jay Koyner, MD (University of Chicago) found that a simple test performed with a one-time dose of the FDAapproved medication furosemide, along with a measurement of urine output, could effectively make this early diagnostic determination. Furosemide is a diuretic that is predominantly cleared by the kidneys, and it can be used to assess certain aspects of the kidneys' function. The test is termed the Furosemide Stress Test (FST).

"These findings may pave the way forward to robust clinical diagnostic tools for clinicians who care for patients with AKI," said Dr. Chawla. "In



this standardized protocol, we were able to determine which patients may or may not require dialysis for <u>acute kidney injury</u> 1 to 3 days ahead of time."

Dr. Chawla noted that using the FST to assess the severity of AKI is similar to the model used in patients with cardiac angina, or chest pain, wherein a marker such as troponin is used in conjunction with an exercise stress test. Similarly, when FST was used in conjunction with AKI markers, it improved predictions of outcomes in patients with early AKI.

In an accompanying editorial, T. Clark Powell, MPH and David Warnock, MD (University of Alabama at Birmingham) noted that the study's findings need to be confirmed and broadened to include larger numbers of individuals. "It will be important to see if the predictive power of the FST remains informative for patients who are not in the critical care setting and <u>critically ill patients</u>," they added.

Provided by American Society of Nephrology

Citation: Simple test detects increased risks in patients with acute kidney injury (2015, February 5) retrieved 7 May 2024 from https://medicalxpress.com/news/2015-02-simple-patients-acute-kidney-injury.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.