

Teen girls from rural areas more likely to have undiagnosed asthma, be depressed

February 24 2015

Teen girls who live in rural areas are more likely than their male counterparts to have undiagnosed asthma, and they often are at a higher risk of depression, according to researchers at the Medical College of Georgia at Georgia Regents University.

According to data collected during a three-year trial of an asthma program tailored to teens, the prevalence of asthma - both diagnosed and undiagnosed - was about the same in rural areas as it is in urban settings. But more girls had undiagnosed asthma than boys.

"There's a lot of speculation about why females are more likely to be undiagnosed," says Dr. Jeana Bush, an MCG Allergy and Immunology Fellow. "Maybe it's because boys are more likely to get a sports physical for athletics and they catch it then. Or maybe it's because girls attribute <u>asthma symptoms</u> to something else, like anxiety. That needs further study."

Bush, who is presenting her findings at the American Academy of Allergy, Asthma and Immunology Annual Meeting, Feb. 20-24 in Houston, also found that the overall rate of <u>depression</u> among asthmatic teens living in rural areas was higher than national averages.

Researchers performed depression screenings using the Depression Intensity Scale Circles (DISC) questionnaire on 332 boys and girls with the disease who were living in four rural counties in Georgia. Of that number, 26 percent - or 87 boys and girls - screened positive for



depression. And of that group, 67 were girls.

"The overall percentage of depression is higher than has been shown in the literature for other chronic diseases, including <u>inflammatory bowel</u> <u>disease</u>, diabetes, <u>congenital heart disease</u>, and even among cancer survivors," Bush says. "That was staggering."

Scientists don't yet fully understand the link between asthma and depression, she says, but they do know that depression is a potential barrier to diagnosis and treatment of the disease.

"Adolescents who are depressed may be less likely to talk about their symptoms or may attribute them to something else," Bush says. "And so much of <u>asthma treatment</u> is about self-management - figuring out your symptoms and preventing an attack when you recognize those symptoms. If you're depressed, you are less likely to be aware of and have the ability to interpret those symptoms."

The data Bush used for her analysis was collected during a National Heart, Lung, and Blood Institute-funded trial of Puff City, led by Dr. Martha Tingen, Charles W. Linder Endowed Chair in Pediatrics, and Dr. Dennis Ownby, Professor of Pediatrics. The program was aimed at helping teens who live in rural areas manage their disease and avoid potentially fatal complications.

Teens from rural areas face a number of problems that could complicate their asthma, including poor housing quality, air pollution, trouble getting to doctors, smaller, less-equipped hospitals, and more exposure to tobacco. Previous studies have shown smoking is more prevalent in rural areas than inner-cities.

Puff City, which featured a "hip" character known as DJ Puffman, approached the issue from three key points - reduction of tobacco



exposure, adherence to medication, and attack readiness.

From 2010-13, data was collected from 2,523 adolescents.

Asthma, which affects more than 300 million people worldwide, is a chronic inflammation that causes a temporary narrowing of the airways that carry oxygen to the lungs. Those narrowings - commonly called <u>asthma</u> attacks - cause more than 4,000 deaths each year in the United States alone, according to the American Lung Association.

Provided by Medical College of Georgia

Citation: Teen girls from rural areas more likely to have undiagnosed asthma, be depressed (2015, February 24) retrieved 6 May 2024 from <u>https://medicalxpress.com/news/2015-02-teen-girls-rural-areas-undiagnosed.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.