

AAD: Telangiectatic photoaging distinct from rosacea subtype

March 23 2015



(HealthDay)—Telangiectatic photoaging (TP) is a distinct dermatological disorder from erythematotelangiectatic subtype of rosacea (ETR), according to a study published online March 23 in *JAMA Dermatology*. The research was published to coincide with the annual meeting of the American Academy of Dermatology, held from March 20 to 24 in San Francisco.

Yolanda R. Helfrich, M.D., from the University of Michigan Medical School in Ann Arbor, and colleagues compared the clinical, histological, and [gene expression](#) features of 26 participants with ETR, 20 with TP, and 11 age- and sex-matched controls.

The researchers identified greater transient erythema in the ETR group

(38 percent graded moderate to severe) versus the TP (0 percent; P

"These data demonstrate that TP is a distinct clinical entity from ETR that can be distinguished on the basis of clinical, histologic, and gene expression findings," the authors write.

Two authors disclosed financial ties to the pharmaceutical industry.

More information: [Abstract](#)

[Full Text](#)

[Editorial](#)

[More Information](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: AAD: Telangiectatic photoaging distinct from rosacea subtype (2015, March 23)
retrieved 4 May 2024 from

<https://medicalxpress.com/news/2015-03-aad-telangiectatic-photoaging-distinct-rosacea.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--