

# Study offers clues to early detection of bipolar disorders in high-risk children

March 3 2015

---



Bipolar disorder is characterized by transitions between depression and mania.  
Credit: Wikipedia

New research published today in the *American Journal of Psychiatry*

indicates a strong link between subthreshold manic episodes and likelihood of developing bipolar disorder in children of parents with bipolar disorder. The study's findings could improve clinical assessment and care for these high-risk children by potentially enabling earlier identification, treatment or possible preventive measures.

The study is among the first to show that subthreshold manic episodes—experiences that approach but do not meet the cut-offs for full-blown [bipolar disorder](#)—are important diagnostic risk factors for the development of bipolar disorder and other mental health conditions in high-risk [children](#). In addition, it found that children of parents with bipolar disorder have substantially higher rates of subthreshold mania or hypomania (13.3 vs 1.2 percent); manic, mixed or hypomanic episodes (9.2 vs 0.8 percent); and major depressive episodes (32.0 vs 14.9 percent) than community children.

"Mental health practitioners should carefully assess for short episodes of manic symptoms, major depression and disruptive behavior disorders in high-risk children, as these are risk factors for the future onset of bipolar disorder," said David Axelson, MD, medical director of Behavioral Health at Nationwide Children's Hospital and senior author on the study. "This may be particularly important if medication is going to be used for treating mood or behavioral symptoms in the child, because some medications might increase the risk of developing mania or psychotic symptoms in someone who is already at very high risk for developing bipolar disorder. Prescribers might choose different medication options, or use lower doses and monitor for problems more closely."

The cumulative rate of bipolar spectrum disorders at age 21 was 23 percent in the high-risk group compared to 3.2 percent in the comparison group. The mean age of onset for a bipolar spectrum disorder was 12 years, with the earliest manic episode being documented at 8 years, highlighting the importance of monitoring subthreshold

symptoms for early intervention.

"Most children who have a parent with bipolar disorder will have a diagnosable psychiatric disorder, such as attention-deficit disorder, oppositional defiant disorder, an anxiety disorder or depression, at some point during their youth," said Dr. Axelson, who is also a professor of clinical psychiatry at The Ohio State University. "However, most children who have a parent with bipolar disorder do not develop bipolar disorder."

Among the study's high-risk children who did develop bipolar disorder, nearly all had identifiable mood episodes and other psychiatric disorders before the onset of full-fledged bipolar illness.

"Short, distinct episodes of manic symptoms were the most specific predictor of developing bipolar disorder in high-risk offspring," Dr. Axelson said. "But so far, many children with these short manic episodes have not progressed to full-fledged bipolar disorder."

This study is part of the Pittsburgh Bipolar Offspring Study (BIOS), which is the largest study to date of children of parents with bipolar disorder. Unique to BIOS is the assessment of children for subthreshold episodes of manic symptoms. This latest publication confirms and extends the baseline findings and describes how mood episodes develop in an effort to aid in early diagnosis on the basis of those subthreshold episodes.

The longitudinal study followed high-risk children and demographically matched offspring of community parents. Follow-up interviews were completed in 91 percent of the offspring with a mean follow-up duration of 6.8 years. The initial and follow-up interviews were conducted using the Structured Clinical Interview for DSM-IV (parents) and the Schedule for Affective Disorders and Schizophrenia for School-Age Children

(children). Follow-ups focused on the assessment of the interval since the previous interview.

"I am continuing to collaborate with researchers on the Pittsburgh Bipolar Offspring Study," Dr. Axelson said. "We expect to find more exciting results that will impact the diagnosis and treatment of high-risk children as we follow the participants into young adulthood."

**More information:** Axelson D, Goldstein B, Goldstein T, Monk K, Yu H, Hickey MB, Sakolsky D, Diler R, Hafeman D, Merranko J, Iyengar S, Brent D, Kupfer D, Birmaher B. Diagnostic precursors to bipolar disorder in offspring of parents with bipolar disorder: A longitudinal study. *American Journal of Psychiatry*. 2015 Mar. [DOI: 10.1176/appi.ajp.2014.14010035](https://doi.org/10.1176/appi.ajp.2014.14010035)

Provided by Nationwide Children's Hospital

Citation: Study offers clues to early detection of bipolar disorders in high-risk children (2015, March 3) retrieved 1 May 2024 from <https://medicalxpress.com/news/2015-03-clues-early-bipolar-disorders-high-risk.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---