

Research explains the formation of long-term motor memory

March 26 2015




Diagram of the cerebellar network. The elements depicted are MF: mossy fibre; GR: granule cells; MLI: molecular layer interneurons; PC: Purkinje cell; VN: vestibular nuclear neuron; and CF: climbing fibre. The weights w and v at PF-PC and MF-VN synapses respectively were modelled to change with time, whereas the weight w_{MLI} at PF-MLI synapses was set to be a constant.

Recent studies of long-term motor memory have pointed out the

involvement of synaptic plasticity at multiple sites in the cerebellum, but the physiological mechanism remains unclear. Now results from a collaboration of researchers at the University of Electro-Communications and the RIKEN Brain Science Institute in Japan, and the University of California, San Diego, in the US, successfully integrated the multiple plasticity mechanisms to explain the formation of long-term motor memory using simulations based on a mathematical model.

Until recently [motor memory](#) was widely considered the result of 'long-term depression' - a state of reduced efficacy following a stimulus - in the synapses at output neurons in the cerebellum called Purkinje cells. Yamazaki and colleagues developed a model for the optokinetic response (OKR) in eye movement that incorporated long-term potentiation - a state of increased efficacy following a stimulus - in the synapses at vestibular nuclear neurons. Thus, the model incorporates two distinct plasticity sites that function synergistically.

The model accurately reproduced experimental results. It showed that an hour's training resulted in a short-term increase of OKR gain, for which long-term depression at the Purkinje cells are responsible. Repetition of this training once a day gradually increased the level of OKR gain after training rather than during it, for which long-term potentiation at the vestibular nucleus neurons are responsible.

"It thus appears as if short-term memory formed in the Purkinje cells during 1-hour training is transferred to the vestibular nuclear neuron after training to consolidate as [long-term memory](#)," concluded the researchers. Their model also reproduced characteristics of the OKR behaviour observed in genetically manipulated mice.

More information: Yamazaki T, Nagao S, Lennon W and Tanaka S
"Modeling memory consolidation during posttraining periods in

cerebellovestibular learning." *PNAS* 112(11): 3541-3546 (2015). [DOI: 10.1073/pnas.1413798112](https://doi.org/10.1073/pnas.1413798112)

Provided by University of Electro-Communications

Citation: Research explains the formation of long-term motor memory (2015, March 26)
retrieved 4 May 2024 from
<https://medicalxpress.com/news/2015-03-formation-long-term-motor-memory.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--