

Improved survival for patients with brain mets who are 50 and younger and receive SRS alone

March 2 2015

Cancer patients with limited brain metastases (one to four tumors) who are 50 years old and younger should receive stereotactic radiosurgery (SRS) without whole brain radiation therapy (WBRT), according to a study available online, open-access, and published in the March 15, 2015 issue of the *International Journal of Radiation Oncology * Biology * Physics (Red Journal)*, the official scientific journal of the American Society for Radiation Oncology (ASTRO). For patients 50 years old and younger who received SRS alone, survival was improved by 13 percentage points when compared to those patients 50 years old and younger who received both SRS and WBRT.

This study, "Phase 3 Trials of Stereotactic Radiation Surgery With or Without Whole-Brain Radiation Therapy For 1 to 4 Brain Metastases: Individual Patient Data Meta-Analysis," analyzed patient data from the three largest randomized clinical trials (RCT) of SRS and WBRT conducted to-date: the Asian trial (JROSG99-1) by Aoyama et al.[1], published in 2006; the North American trial (MDACC NCT00548756) by Chang et al.[2], published in 2009; and the European trial (EORTC 22952-26001) by Kocher et al.[3], published in 2011. A total of 364 patients from the three RCTs were evaluated for this meta-analysis. Of those 364 patients, 51 percent (186) were treated with SRS alone, and 49 percent (178) received both SRS and WBRT. Nineteen percent of patients (68) were 50 years old and younger, and 61 percent (19) of these patients had a single brain metastasis. Twenty percent of all patients (72)

had local brain failure, which is the occurrence of progression of previously treated brain metastases; and 43 percent (156) experienced distant brain failure, which is the occurrence of new brain metastases in areas of the brain outside the primary tumor site(s).

The impact of age on treatment effectiveness revealed SRS alone yielded improved overall survival (OS) in patients 50 years old and younger. Patients 50 years old and younger who received SRS alone had a median survival of 13.6 months after treatment, a 65 percent improvement, as opposed to 8.2 months for patients 50 years old and younger who were treated with SRS plus WBRT. Patients >50 years old had a median survival of 10.1 months when treated with SRS alone, and 8.6 months for those who received SRS plus WBRT.

"We expected to see a survival advantage favoring combined therapy of SRS and WBRT. However, these data clearly demonstrate the benefit for SRS alone to improve survival for our younger patients with limited brain metastases," said lead author of the study Arjun Sahgal, MD, associate professor of [radiation oncology](#) and surgery at the University of Toronto, and a radiation oncologist at the Odette Cancer Centre of the Sunnybrook Health Sciences Centre in Toronto. "Furthermore, it was previously thought that the positive effect of whole brain radiation in reducing the risk of distant brain relapse was generalizable for all patients. However, we did not observe this effect in patients 50 years and younger with limited [brain metastases](#). In these patients, the same rate of distant brain failure was observed despite treatment with whole brain radiation. This result, together with our survival result, gave rise to the hypothesis that if patients are treated with whole [brain radiation](#) without realizing the benefits of improving distant brain control, then survival may be adversely affected. Therefore, our sub-group meta-analysis has swung the pendulum in favor of SRS alone as the standard of care. These findings also reinforce ASTRO's Choosing Wisely recommendation[4] that states that it may not be necessary to add WBRT to SRS, thus

improving [patients'](#) quality of life and memory function."

More information: Drs. Nils D. Arvold and Paul J. Catalano have reviewed Sahgal et al.'s research. Their editorial, "Local Therapies for Brain Metastases, Competing Risks, and Overall Survival," is also published in the March 15, 2015, issue of the *Red Journal*.

[1] Aoyama H, Shirato H, Tago M, et al. Stereotactic radiosurgery plus whole-brain radiation therapy vs stereotactic radiosurgery alone for treatment of bone metastases: A randomized controlled trial. *JAMA* 2006;295:2483-2491.

[2] Chang EL, Wefel JS, Hess KR, et al. Neurocognition in patients with brain metastases treated with radiosurgery or radiosurgery plus whole-brain irradiation: a randomised controlled trial. *Lancet Oncol* 2009;10:1037-1044.

[3] Kocher M, Soffietti R, Abacioglu U, et al. Adjuvant whole-brain radiotherapy versus observation after radiosurgery or surgical resection of one to three cerebral metastases: results of the EORTC 22952-26001 study. *J Clin Oncol* 2010;29:134-141.

[4] ASTRO's Choosing Wisely List. ABIM Foundation.
[www.choosingwisely.org/doctor- ... -radiation-oncology/](http://www.choosingwisely.org/doctor-...-radiation-oncology/)

Provided by American Society for Radiation Oncology

Citation: Improved survival for patients with brain mets who are 50 and younger and receive SRS alone (2015, March 2) retrieved 6 May 2024 from
<https://medicalxpress.com/news/2015-03-survival-patients-brain-mets-younger.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.