

Youth suicide rate in rural areas is nearly double the rate in cities

March 9 2015

The adolescent and young-adult suicide rate in the United States was almost twice as high in rural settings than in urban areas between 1996 and 2010, and new research suggests that the gap appears to be widening.

Of the nearly 67,000 suicides analyzed, the rate of suicide for both males and females living in rural areas was about double the rate in cities.

The research, published in *JAMA Pediatrics*, also showed that gun use has decreased and that hanging has become a more common method of youth suicide for both males and females. Suicide rates by firearm and hanging were both disproportionately higher in rural areas than in urban regions for both sexes.

The findings suggest there is an urgent need to improve access to and availability of mental health care in rural areas, researchers say. The authors suggest that incorporating mental health care into primary care settings, providing care via telemedicine and creating school-based interventions might help reduce youth suicides in rural areas.

"These kinds of surveillance studies can really help us identify areas to target our prevention efforts, and it's clear we need to target rural areas for primary prevention of suicide," said Cynthia Fontanella, clinical assistant professor of psychiatry and behavioral health at The Ohio State University Wexner Medical Center and lead author of the study.

Fontanella cited access barriers to health care, geographic isolation and disproportionate stigma associated with mental illness as potential reasons for the observed disparities. "Rural America is beautiful, and many individuals live in close-knit communities. But in terms of access to services and perhaps some other factors, the cards may be stacked against them."

Fontanella and colleagues analyzed mortality data from the National Center for Health Statistics National Vital Statistics System. Between 1996 and 2010, 66,595 youths between age 10 and 24 died by suicide in the United States.

Among males, the rural versus urban suicide rate was 19.93 and 10.31 per 100,000, respectively, and for females, 4.40 and 2.39 per 100,000, respectively. When the researchers controlled for a number of other potential contributing factors, the rural and urban suicide rate disparity increased over time for males. The analysis also showed the suicide rate for young males is four times higher than for young females.

More than half of this population - 51.1 percent - committed suicide with a firearm, and 33.9 percent died by hanging. Poisoning accounted for 7.9 percent of suicides and other methods, such as jumping and transportation-related suicides, composed 7.1 percent of the deaths.

Clinicians have identified many risk factors for suicide over the years, but two in particular stand out, said John Campo, chair of psychiatry and behavioral health at Wexner Medical Center and senior author of the paper. About 90 percent of people who commit suicide suffer from a major psychiatric problem such as depression or bipolar disorder. Substance abuse, which can increase impulsivity and interfere with mood, is also frequently linked to suicides.

No studies have been done to try to pinpoint specific causes of this

disparity in suicides based on where young people live. But these authors considered a number of potential contributing factors in the paper, all supported by previous research.

Between the dispersed nature of rural populations and a severe shortage of mental health practitioners, access to care is a major problem.

"If a rural child is depressed, it's much harder to get state-of-the-art care. And it's especially difficult to receive psychotherapy in a rural area," Campo said.

Of the 1,669 federally designated shortage areas for mental health services in the United States, 85 percent are in rural regions. And more than half of the counties in this country - all rural - do not have a single psychiatrist, psychologist or social worker serving the region.

This lack of access can compound the severity of illness, Fontanella said. People living in rural areas have to travel longer distances to see health care practitioners and wait longer for appointments, so by the time they receive care they may have developed more serious symptoms. And the culture of self-reliance in rural areas, as compared to urban living, may intensify the stigma associated with mental illness.

"There is a lot of pride around taking care of your own problems and turning to neighbors rather than professionals for help," Fontanella said.

Because families living in rural areas tend to see a primary care practitioner for most of their health care, integrating mental health care into rural primary care settings - either with on-site practitioners or via remote video access - has promise, Campo said.

"It's going to be hard to generate the numbers of mental-health providers to get where we need to go. So we need to address things qualitatively to

improve the entire system of care, which will improve access," he said.

More information: *JAMA Pediatr.* Published online March 9, 2015.
[DOI: 10.1001/jamapediatrics.2014.3561](https://doi.org/10.1001/jamapediatrics.2014.3561)

JAMA Pediatr. Published online March 9, 2015. [DOI: 10.1001/jamapediatrics.2015.104](https://doi.org/10.1001/jamapediatrics.2015.104)

Provided by The Ohio State University

Citation: Youth suicide rate in rural areas is nearly double the rate in cities (2015, March 9)
retrieved 6 May 2024 from
<https://medicalxpress.com/news/2015-03-widening-rural-urban-disparities-youth-suicides.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|