

# Babies feel pain 'like adults'

April 21 2015

---



Credit: Anna Langova/public domain

The brains of babies 'light up' in a very similar way to adults when exposed to the same painful stimulus, a pioneering Oxford University brain scanning study has discovered. It suggests that babies experience pain much like adults.

The study looked at 10 healthy infants aged between one and six days old and 10 healthy [adults](#) aged 23-36 years. Infants were recruited from the John Radcliffe Hospital, Oxford (UK) and adult volunteers were

Oxford University staff or students.

During the research [babies](#), accompanied by parents and clinical staff, were placed in a Magnetic Resonance Imaging (MRI) scanner where they usually fell asleep. MRI scans were then taken of the babies' brains as they were 'poked' on the bottom of their feet with a special retracting rod creating a sensation 'like being poked with a pencil' - mild enough that it did not wake them up. These scans were then compared with brain scans of adults exposed to the same pain stimulus.

The researchers found that 18 of the 20 brain regions active in adults experiencing pain were active in babies. Scans also showed that babies' brains had the same response to a weak 'poke' (of force 128mN) as adults did to a stimulus four times as strong (512mN). The findings suggest that not only do babies experience pain much like adults but that they also have a much lower pain threshold.

The research was funded by the Wellcome Trust and is reported in the journal *eLife*.

'Up until recently people didn't think it was possible to study pain in babies using MRI because, unlike adults, they don't keep still in the scanner!' said Dr Rebecca Slater of Oxford University's Department of Paediatrics, lead author of the report. 'However, as babies that are less than a week old are more docile than older babies, we found that their parents were able to get them to fall asleep inside a scanner so that, for the first time, we could study pain in the infant brain using MRI.

'This is particularly important when it comes to pain: obviously babies can't tell us about their experience of pain and it is difficult to infer pain from visual observations. In fact some people have argued that babies' brains are not developed enough for them to really 'feel' pain, any reaction being just a reflex - our study provides the first really strong

evidence that this is not the case.'

The researchers say that it is now possible to see pain 'happening' inside the infant brain and it looks a lot like pain in adults.

As recently as the 1980s it was common practice for babies to be given neuromuscular blocks but no [pain relief](#) medication during surgery [1]. In 2014 a review of neonatal [pain management](#) practice in intensive care highlighted that although such infants experience an average of 11 painful procedures per day 60% of babies do not receive any kind of pain medication [2].

'Thousands of babies across the UK undergo painful procedures every day but there are often no local pain management guidelines to help clinicians. Our study suggests that not only do babies experience pain but they may be more sensitive to it than adults,' said Dr Slater. 'We have to think that if we would provide pain relief for an older child undergoing a procedure then we should look at giving pain relief to an infant undergoing a similar procedure.'

Dr Slater added: 'Recent studies in adults have shown that it is possible to detect a neurological signature of pain using MRI. In the future we hope to develop similar systems to detect the 'pain signature' in babies' brains: this could enable us to test different pain relief treatments and see what would be most effective for this vulnerable population who can't speak for themselves.'

Rachel Edwards, who is 33 and from Oxford, gave permission for her son Alex to take part in the study. Alex was the first baby to be placed in the MRI scanner:

Rachel Edwards said: 'People know so little about how babies feel [pain](#), you can tell they are in distress from their reaction and I was curious

about why they react in the way they do.

'Before Alex went in I got to feel all the things he would feel as part of the study including the pencil-like retracting rod: it wasn't particularly painful, it was more of a precise feeling of touch.

'It was really reassuring how highly skilled the staff were and how good they were at calming and handling him.

'I fed him and put him down on this special bean bag that they suck all the air out of to help keep the babies' heads still. Then they put on earphones that cut out some of the sound but I think the noise actually soothed him. He was out for the count, he didn't wake up during the scanning and seemed really content afterwards.'

**More information:** A report of the research is published in the journal *eLife*: [dx.doi.org/10.7554/eLife.06356](https://doi.org/10.7554/eLife.06356)

Notes:

[1] Anand KJ, and Hickey PR. Pain and its effects in the human neonate and fetus. *The New England journal of medicine*. 1987;317(21):1321-9.

[2] Roofthoof DW, Simons SH, Anand KJ, Tibboel D, and van Dijk M. Eight years later, are we still hurting newborn infants? *Neonatology*. 2014;105(3):218-26.

Provided by Oxford University

Citation: Babies feel pain 'like adults' (2015, April 21) retrieved 5 May 2024 from <https://medicalxpress.com/news/2015-04-babies-pain-adults.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.