

Oophorectomy associated with decrease in breast cancer death in women with cancer, BRCA1 mutation

April 23 2015

Removal of the ovaries, a procedure known as an oophorectomy, was associated with a 62 percent reduction in breast cancer death in women diagnosed with breast cancer and carrying a BRCA1 gene mutation, according to an article published online by *JAMA Oncology*.

Women who carry a germline mutation in either the BRCA1 or BRCA2 gene face a lifetime risk of [breast cancer](#) of up to 70 percent. Once they are diagnosed with breast cancer, they face high risks of both second primary breast and ovarian cancers. Other studies of BRCA gene mutation carriers have reported reduced mortality associated with oophorectomy for women with a history of breast cancer, according to the study background.

Steven A. Narod, M.D., and Kelly Metcalfe, Ph.D., of the Women's College Research Institute, Toronto, Canada, and coauthors sought to confirm these earlier observations in a group of women with BRCA1 and BRCA2 gene mutations and early-stage breast cancer. Their study included 676 women, of whom 345 underwent oophorectomy after being diagnosed with breast cancer, while 331 women retained both ovaries.

The study found 20-year survival for the entire group was 77.4 percent. In the entire group, there was a 56 percent reduction in breast [cancer death](#) associated with oophorectomy. Undergoing an oophorectomy was

associated with a significant reduction (62 percent) in breast cancer death in women with a BRCA1 mutation but not in women with a BRCA2 mutation because the 43 percent reduction authors found was not statistically significant.

In addition there were nine deaths from ovarian cancer in the group of women who did not have oophorectomies. The authors found a 65 percent reduction in all-cause mortality associated with oophorectomy in their analysis.

According to the study results, oophorectomies were performed an average of six years after breast cancer diagnosis. For the 70 BRCA1 carriers for whom the oophorectomy was performed within two years of breast cancer diagnosis, there was a 73 percent reduction in death compared with women with a BRCA1 mutation who never underwent oophorectomy. The authors note the protective effect of oophorectomy on deaths from breast cancer was apparent immediately after diagnosis and lasted for 15 years.

"It is important that follow-up studies be performed on women who undergo oophorectomy as part of their initial treatment, in particular, those women who undergo oophorectomy in the first year after diagnosis. It is also important that our observations be confirmed in other study populations. Further data are needed, in particular for BRCA2 carriers in order to confirm the benefit of oophorectomy in this population," article concludes.

In a related editor's note, Mary L. Disis, M.D., editor-in-chief of *JAMA Oncology*, writes: "The results provide a validation of the role of oophorectomy in conveying both a disease-free and overall survival benefit for BRCA1 mutation carriers. Oophorectomy after the primary diagnosis of breast cancer significantly reduced breast cancer-specific mortality in women with BRCA1 mutations but not in BRCA2 mutation

carriers. In the entire group, oophorectomy was particularly effective for survival benefit in [women](#) with estrogen receptor-negative breast cancer. ... The data reported here are compelling and suggest that the potential of oophorectomy should become part of the treatment discussion at the time of [diagnosis](#) for BRCA mutation carriers with early-stage breast cancers."

More information: *JAMA Oncol.* Published online April 16, 2015.
[DOI: 10.1001/jamaoncol.2015.0658](https://doi.org/10.1001/jamaoncol.2015.0658)

Provided by The JAMA Network Journals

Citation: Oophorectomy associated with decrease in breast cancer death in women with cancer, BRCA1 mutation (2015, April 23) retrieved 5 May 2024 from
<https://medicalxpress.com/news/2015-04-oophorectomy-decrease-breast-cancer-death.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--