


Women show persistent memory impairment after concussion

April 28 2015



Working memory functional activation in healthy control subjects and patients with MTBI is shown. Surface-rendered projection on a representative atlas brain shows areas of activation in response to increased working memory load. Note the presence of increased activation in bilateral frontal and parietal regions, a finding consistent with activation of working memory circuitry in each group. A: In control subjects, visual comparison of working memory activation patterns shows more activation in women than in men, especially in the frontal region. B: There is substantially less activation seen in female patients with MTBI than in female control subjects, and there is substantially more activation seen in male patients with MTBI than in male control subjects. C: At the 6-week follow-up study, female patients with MTBI showed a persistent hypoactivation pattern,

whereas male patients with MTBI showed regression of hyperactivation and demonstrated a return to similar activation levels as those in male control subjects. Credit: Radiological Society of North America

Women may have a more difficult time than men in recovering from concussion, according to a new study published online in the journal *Radiology*.

Concussion, also known as mild [traumatic brain injury](#) (MTBI), is a common medical problem affecting cognitive function and quality of life in some individuals.


In most cases, patients who experience MTBI will recover fully, typically within three months. Ten to 15 percent of MTBI patients will continue to experience persistent disabling problems beyond three months, which can include post-traumatic headache, sleep disturbance, loss of balance, memory and other cognitive impairments, fatigue, and mood or affective disorders.

The lasting effects of concussion have received widespread attention in recent years and caused growing concern, particularly among amateur and professional athletes and sports organizations. Several reports have indicated that female athletes suffer concussions at a higher rate than male athletes playing similar sports.

"In clinical practice, more women than men seek medical attention due to persistent symptoms after MTBI at a ratio of almost 2:1," said the study's lead author, Chi-Jen Chen, M.D., from the Department of Diagnostic Radiology and Brain and Consciousness Research Center at Taipei Medical University Shuang-Ho Hospital and superintendent at Chia-Yi Hospital in Taiwan. "We started to wonder whether there might

be differences in MTBI outcomes between men and women."

Dr. Chen and colleagues set out to objectively evaluate gender differences in MTBI by using functional magnetic resonance imaging (fMRI) to analyze [brain activation patterns](#) during [working memory](#) tasks.




Sex differences of working memory functional activation in healthy control subjects in response to increased working memory load. A voxel-by-voxel comparison of female (left image) and male (right image) control subjects shows more activation (arrows) in female control subjects than in male control subjects. Credit: Radiological Society of North America

"Difficulty with working memory is a commonly reported impairment after MTBI," Dr. Chen said. "Since working memory is important for a wide variety of cognitive skills, compromised working memory could have significant effects on everyday life."

The researchers studied 30 patients with MTBI and 30 control patients. Both groups contained an equal number of men and women. Patients underwent an fMRI exam within one month after injury and a follow-up fMRI exam at six weeks after first scan. All participants underwent neuropsychological tests, including digit span and continuous performance test (CPT). Digit span is a short-term memory test measuring how many numbers a participant can remember in sequence. CPT measures a person's sustained and selective attention and impulsivity.

Initial fMRI results of the MTBI patients showed increased activation in working memory brain circuits in the men and decreased activation in the women, compared to the controls. At follow-up, men with MTBI returned to a normal activation pattern similar to the controls, whereas the women showed persistent hypoactivation, suggesting ongoing working memory problems.



Sex differences of working memory functional activation at initial study of patients with MTBI compared with control subjects in response to increased working memory load. Voxel-by-voxel comparison of initial studies of patients with MTBI and control subjects shows a little more activation in the working memory circuitry (arrowheads) in female control subjects than in female patients with MTBI. However, substantially more activation (arrows) is seen in male patients with MTBI than in male control subjects. Credit: Radiological Society of North America

Neuropsychological results showed that among the women, the total digit

span score was lower in the MTBI group, compared to the control group.

"These findings provide evidence that female gender may be a risk factor for working memory impairment after MTBI," Dr. Chen said. "If so, more aggressive management should be initiated once MTBI is diagnosed in female patients."

Because fMRI has the potential to detect working memory impairment, predict outcome and monitor treatment effect, incorporation of fMRI into treatment protocol for MTBI may be advisable in the future, according to Dr. Chen. However, he cautions that further study needs to be done to validate these preliminary findings.

More information: Sex Differences in Working Memory after Mild Traumatic Brain Injury: A Functional MR Imaging study, *Radiology*, 2015.

Provided by Radiological Society of North America

Citation: Women show persistent memory impairment after concussion (2015, April 28) retrieved 11 May 2024 from <https://medicalxpress.com/news/2015-04-women-persistent-memory-impairment-concussion.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--