

## Autism and rare childhood speech disorder often coincide

May 27 2015

---



Quinn, an autistic boy, and the line of toys he made before falling asleep. Repeatedly stacking or lining up objects is a behavior commonly associated with autism. Credit: Wikipedia.

Some children with autism should undergo ongoing screenings for apraxia, a rare neurological speech disorder, because the two conditions often go hand-in-hand, according to Penn State College of Medicine researchers.

Over the course of a three-year study, 64 percent of [children](#) initially diagnosed with [autism](#) were found to also have apraxia. The study also showed that the commonly used Checklist for Autism Spectrum Disorder (CASD) accurately diagnoses autism in children with apraxia.

"Children with apraxia have difficulty coordinating the use of their tongue, lips, mouth and jaw to accurately produce speech sounds, so that each time they say the same word, it comes out differently, and even their parents have difficulty understanding them," said Cheryl Tierney, associate professor of pediatrics.

It's estimated that one in 68 children in the United States has autism and one to two in 1,000 have apraxia. With increased recognition and improved evaluation measures, more children are being identified with autism and apraxia. Developmental experts have long noted autism and apraxia seem to frequently coincide.

The researchers studied 30 children between the ages of 15 months and five years who were referred to the Penn State Hershey Pediatric Developmental Communication Assessment Clinic for concerns regarding speech, language or autism. After an initial diagnostic screening, follow-up testing showed that 63.6 percent of children initially diagnosed with autism also had apraxia and 36.8 percent of children initially diagnosed with apraxia also had autism.

The researchers determined that about a quarter of children studied—23.3 percent—had both autism and apraxia at the initial diagnosis, and the same amount had neither condition. Researchers report their results in the *Journal of Developmental and Behavioral Pediatrics*.

Outcomes for both autism and apraxia can be improved with early intervention, so the findings point to the importance of ongoing apraxia

assessment for some children with autism, and vice versa.

Based on the study, Tierney believes that children with autism or apraxia who are non-verbal should be screened for the other condition until they start talking.

The new study also verified that the CASD, a commonly used autism tool, does not over-diagnose autism in children with apraxia.

Penn State College of Medicine psychologist Susan Mayes developed the CASD. It was standardized on children with language disorders—trouble in using language for expression—but not for speech disorders—trouble producing the sound—so clinicians have wondered if it could accurately be used in children with apraxia.

"We were concerned that children with childhood apraxia of speech were being mislabeled as having autism when they were being evaluated for it, but we found that this was not the case," Tierney said.

To test the accuracy of the CASD in children with apraxia, the researchers used four other tools, which included two assessments for autism and two assessments for apraxia. Overall, the CASD was 97 percent sensitive, and it accurately diagnosed autism in 95 percent of children with apraxia.

"We found that in our diagnostic center, we were not over-diagnosing autism in a cohort of children that had a concern for childhood apraxia of speech," Tierney said.

Autism and apraxia have very different interventions, so making a correct diagnosis is important for preventing long-term problems.

"CASD is used to diagnose autism, but it can also be used to rule out

autism," Tierney said. "We wanted to make sure this tool came out negative in a child that just had apraxia and we found that it really does help us distinguish kids accurately. Children who had apraxia without autism, children who had autism but not [apraxia](#), or those who had both or neither were picked up accurately. The tool was very sensitive in picking up all those different combinations. And that's good news because we want the right treatment for the right child for the right diagnosis."

Provided by Pennsylvania State University

Citation: Autism and rare childhood speech disorder often coincide (2015, May 27) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2015-05-autism-rare-childhood-speech-disorder.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--