

## **Diagnosing cancer with help from bacteria**

May 27 2015



Electron micrograph image of a cluster of E. coli bacteria magnified 10,000 times. Researchers were able to quickly sequence the DNA of a suspected strain of E. coli thought to be the case of the latest outbreak in Europe. Credit: Eric Erbe | Christopher Pooley | USDA | ARS | EMU

Engineers at MIT and the University of California at San Diego (UCSD) have devised a new way to detect cancer that has spread to the liver, by enlisting help from probiotics—beneficial bacteria similar to those found in yogurt.

Many types of cancer, including colon and pancreatic, tend to metastasize to the liver. The earlier doctors can find these tumors, the more likely that they can successfully treat them.

"There are interventions, like local surgery or local ablation, that physicians can perform if the spread of disease in the liver is confined,



and because the liver can regenerate, these interventions are tolerated. New data are showing that those patients have a higher survival rate, so there's a particular need for detecting early metastasis in the liver," says Sangeeta Bhatia, the John and Dorothy Wilson Professor of Health Sciences and Electrical Engineering and Computer Science at MIT.

Using a harmless strain of *E. coli* that colonizes the liver, the researchers programmed the bacteria to produce a luminescent signal that can be detected with a simple urine test. Bhatia and Jeff Hasty, a professor of bioengineering at UCSD, are the senior authors of a paper describing the new approach this week in the journal *Science Translational Medicine*. Lead authors are MIT postdoc Tal Danino and UCSD postdoc Arthur Prindle.

## **Microbial help**

Previous studies had shown that bacteria can penetrate and grow in the tumor microenvironment, where there are lots of nutrients and the body's immune system is compromised. Because of this, scientists have been trying for many years to develop bacteria as a possible vehicle for cancer treatment.

The MIT and UCSD researchers began exploring this idea a few years ago, but soon expanded their efforts to include the concept of creating a bacterial diagnostic.

To turn bacteria into diagnostic devices, the researchers engineered the cells to express the gene for a naturally occurring enzyme called lacZ that cleaves lactose into glucose and galactose. In this case, lacZ acts on a molecule injected into the mice, consisting of galactose linked to luciferin, a luminescent protein naturally produced by fireflies. Luciferin is cleaved from galactose and excreted in the urine, where it can easily be detected using a common laboratory test.



At first, the researchers were interested in developing these bacteria for injection into patients, but then decided to investigate the possibility of delivering the bacteria orally, just like the probiotic bacteria found in yogurt. To achieve that, they integrated their diagnostic circuits into a harmless strain of *E. coli* called Nissle 1917, which is marketed as a promoter of gastrointestinal health.

In tests with mice, the researchers found that orally delivered bacteria do not accumulate in tumors all over the body, but they do predictably zero in on liver tumors because the hepatic portal vein carries them from the digestive tract to the liver.

"We realized that if we gave a probiotic, we weren't going to be able to get bacteria concentrations high enough to colonize the tumors all over the body, but we hypothesized that if we had tumors in the liver they would get the highest dose from an oral delivery," says Bhatia, who is a member of MIT's Koch Institute for Integrative Cancer Research and Institute for Medical Engineering and Science.

This allowed the team to develop a diagnostic specialized for liver tumors. In tests in mice with colon cancer that has spread to the liver, the <u>probiotic bacteria</u> colonized nearly 90 percent of the metastatic tumors.

In the mouse experiments, animals that were given the engineered bacteria did not exhibit any harmful side effects.

## More sensitive detection

The researchers focused on the liver not only because it is a natural target for these bacteria, but also because the liver is hard to image with conventional imaging techniques like CT scanning or magnetic resonance imaging (MRI), making it difficult to diagnose <u>metastatic</u> <u>tumors</u> there.



With the new system, the researchers can detect <u>liver tumors</u> larger than about one cubic millimeter, offering more sensitivity than existing imaging methods. This kind of diagnostic could be most useful for monitoring patients after they have had a colon tumor removed because they are at risk for recurrence in the <u>liver</u>, Bhatia says.

The MIT team is now pursuing the idea of using probiotic <u>bacteria</u> to treat cancer, as well as for diagnosing it.

For background on this project, view this video presentation:

**More information:** "Programmable probiotics for detection of cancer in urine," *Science Translational Medicine*, <u>stm.sciencemag.org/lookup/doi/ ... scitranslmed.aaa3519</u>

Related paper: "Detection of pathological biomarkers in human clinical samples via amplifying genetic switches and logic gates," <u>stm.sciencemag.org/lookup/doi/ ... scitranslmed.aaa3601</u>

## Provided by Massachusetts Institute of Technology

Citation: Diagnosing cancer with help from bacteria (2015, May 27) retrieved 2 May 2024 from <u>https://medicalxpress.com/news/2015-05-cancer-bacteria.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.