

Study characterizes the incidence and effects of severe kidney injury during pregnancy

May 14 2015

A new study indicates that severe kidney injury is rare during pregnancy, but it typically occurs in otherwise healthy women who acquire a major pregnancy-related complication. The study, which will appear in an upcoming issue of the *Journal of the American Society of Nephrology* (*JASN*), also reveals the health outcomes of mothers and their babies following severe kidney injury during pregnancy.

Acute [kidney injury](#) (AKI), an abrupt or rapid decline in [kidney function](#), is a rare but serious complication of [pregnancy](#). Although previously considered a disappearing entity, the incidence and outcomes of AKI during pregnancy may be adversely impacted as increasing numbers of pregnancies are occurring in women who are older, have conditions such as diabetes or chronic kidney disease, or use reproductive technologies.

To determine the incidence, characteristics, and outcomes of women with AKI who require treatment with dialysis during pregnancy and the postpartum period, Ainslie Hildebrand, MD (Western University's London Health Sciences Centre, in Canada) and her colleagues analyzed information on all pregnancies from 1997 to 2011 in Ontario.

Among the major findings:

- The incidence of AKI that was treated with dialysis was 1 in 10,000 pregnancies (188 out of 1.9 million pregnancies).
- Most women who developed AKI during pregnancy had no

recorded pre-existing health conditions; however, compared with healthy women, those with pre-pregnancy hypertension, diabetes, [chronic kidney disease](#), or lupus were at least twice as likely to develop AKI during pregnancy, and those with a major pregnancy-related complication such as preeclampsia were nearly 4-times more likely to develop AKI.

- Four percent of women who developed AKI during pregnancy died during the 15-year study, and 4% of survivors remained dialysis-dependent after delivery.
- Adverse perinatal outcomes—such as low birth weight and preterm birth—occurred in 35% of pregnancies affected by AKI; however there were no stillbirths and fewer than 5 neonatal deaths (

Citation: Study characterizes the incidence and effects of severe kidney injury during pregnancy (2015, May 14) retrieved 27 April 2024 from <https://medicalxpress.com/news/2015-05-characterizes-incidence-effects-severe-kidney.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.