

# Changes observed in HbA1c during ramadan

May 20 2015

---



(HealthDay)—For patients with type 2 diabetes, during Ramadan, the greatest change among metabolic parameters is seen for glycemia, according to a study published online May 13 in the *Journal of Diabetes Investigation*.

Melanie Y.L. Siaw, from the National University of Singapore, and colleagues conducted a retrospective study using a national electronic database to examine trends in hemoglobin A1c (HbA1c), [systolic blood pressure](#) (SBP), [low-density lipoprotein cholesterol](#) (LDL-C), and triglycerides (TG) in [patients](#) with type 2 diabetes during Ramadan. A total of 5,172 eligible patients were categorized according to their HbA1c control before Ramadan: Group 1, HbA1c  $\geq 10$  percent; Group 2, HbA1c, 7.1 to 9.9 percent; Group 3, HbA1c  $\leq 7.0$  percent.

The researchers observed variation in the mean change of HbA1c, from  $-1.4$  to  $+0.2$  percent, with the greatest decrease seen in Group 1 (P

"In conclusion, among the known metabolic parameters, the greatest change was observed in glycemia during Ramadan," the authors write. "Patients with a mean baseline HbA1c value of 10 percent or more experienced the greatest HbA1c reduction."

**More information:** [Abstract](#)  
[Full Text](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: Changes observed in HbA1c during ramadan (2015, May 20) retrieved 20 March 2024 from <https://medicalxpress.com/news/2015-05-hba1c-ramadan.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--