


Modeling memory in the brain

May 18 2015, by Nik Papageorgiou



Credit: ThinkStock

Scientists at EPFL have uncovered mathematical equations behind the way the brain forms – and even loses – memories.

Memory is one of the most crucial elements of life. Without memory, there is no learning; without learning there is no invention, progress, or civilization. On the flipside, forgetting some experiences, especially traumatic ones, can help regain mental health and function. The key to all this is to understand how the brain forms memories in the first place, and then how it retains and recalls them. Scientists at EPFL have developed a mathematical model to describe how networks of [neurons](#) create memories. Published in *Nature Communications*, the model could clarify longstanding theories of [memory formation](#), and could change the way we understand, simulate and even alter memory formation.

"Fire together, wire together"

Neurons form networks through specialized connections called "synapses". The neuron sending a signal through a synapse is called "pre-synaptic", and the one receiving it "post-synaptic". Synapses show a lot of plasticity, allowing neurons to change their communication speed and intensity. Today, synaptic plasticity is considered the basis of how we learn and make memories.

The most prevalent theory of synaptic plasticity, named after neuroscientist Donald Hebb, states that a synapse becomes stronger when the pre-synaptic neuron fires repeatedly and stimulates the post-synaptic neuron to fire in sync. In such Hebbian synapses, "cells that fire together, wire together." However, models of Hebbian plasticity fail to simulate memory formation accurately because they cannot account for external biological factors, and also because not all synapses are Hebbian.

A realistic model of memory

A research team led by Wolfram Gerstner at EPFL has now developed a

model of Hebbian plasticity that succeeds where previous ones have failed. The researchers focused on the formation of what are known as "memory assemblies", which are networks of neurons, connected via synapses, which can store a particular segment of a memory. When a memory is being recalled, its particular assemblies piece it together to produce a whole.

The researchers used a third-generation neural network model called a "spiking neural network" (SNN). In an SNN, when a neuron fires a signal (a spike), it travels to other neurons, which respond accordingly by increasing or decreasing their own ability to fire a signal, thereby strengthening or weakening the connection.

Different synapses, different timescales

Gerstner's team simulated hundreds of SNNs in order to explore different types of synapses and multiple forms of synaptic plasticity across different timescales. The simulations suggested that memory formation and recall actually follows a "well-orchestrated combination" of both Hebbian and non-Hebbian rules of synaptic plasticity. In other words, the formation of memory assemblies does not depend only on the signals coming from the pre-synaptic neuron but also on input from external neurons that indirectly modulate the strength of the synapse.

Time was also a crucial component of the simulation. Once a memory assembly had been formed, memories could be recalled even days after by selectively triggering the activity of specific neurons in the memory assembly. Overall, the scientists' model suggests that the wide diversity of [synaptic plasticity](#) in the brain (Hebbian, non-Hebbian and even other types) is "orchestrated towards achieving common functional goals."

From their results, the scientists were able to derive a complex algorithm that currently is the most accurate representation of this complex

phenomenon. The algorithm can be adapted to help develop next-level simulations of memory formation and recall, which can advance our understanding of how the brain works as whole. In addition, the findings can inform strategies for addressing with traumatic memories or even improving educational efforts. "If we can understand how [synapses](#) work together to forge or dismantle [memory](#) networks, we can advance fields such as cognition and psychotherapy," says Gerstner.

More information: "Diverse synaptic plasticity mechanisms orchestrated to form and retrieve memories in spiking neural networks." *Nature Communications* 6:6922, 21 April 2015. [DOI: 10.1038/ncomms7922](#)

Provided by Ecole Polytechnique Federale de Lausanne

Citation: Modeling memory in the brain (2015, May 18) retrieved 10 May 2024 from <https://medicalxpress.com/news/2015-05-memory-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
