

80 percent of cervical cancers found to be preventable with latest 9-valent HPV vaccine

May 11 2015

The new 9-valent human papillomavirus vaccine, can potentially prevent 80 percent of cervical cancers in the United States, if given to all 11- or 12-year-old children before they are exposed to the virus.

In addition to protecting against 80 percent of cervical cancers, the new 9-Valent human-papillomavirus-vaccine, which includes seven cancer causing HPV-types - 16,18,31,33,45,52 and 58 - has the potential to protect against nearly 19,000 other cancers diagnosed in the United States, including anal, oropharyngeal and penile cancers. This is a 13 percent increase in protection against HPV-related cancers in comparison to the first vaccines on the market, Gardasil and Cervarix, which protected against HPV types 16 and 18.

These findings come from a seven-center study published in the *Journal* of the National Cancer Institute. The Centers for Disease Control and Prevention initiated the research effort, in conjunction with Cedars-Sinai.

"This is the first comprehensive study of its kind and shows the potential to not only reduce the global cancer burden, but also guide clinical decision-making with regard to childhood vaccinations," said Marc T. Goodman, PhD, MPH, senior author of the study and director of Cancer Prevention and Genetics at the Cedars-Sinai Samuel Oschin Comprehensive Cancer Institute.

The study found the 9-Valent vaccine, under the trademark of



Gardasil-9, also has the potential to protect against an additional 8 percent of oropharyngeal cancers, which include the base of the tongue and tonsils. This disease is the second-most-common HPV-associated cancer.

"We found that 70 percent of patient DNA tissue samples with cancer of the oropharynx harbored HPV," added Goodman. "This is a much higher percentage of HPV than observed in other studies, likely because of changes in sexual behaviors, such as increased oral-genital contact."

The 9-Valent <u>vaccine</u> was also found to potentially increase protection from other HPV-related cancers including those of the vulva, from 71 to 92 percent; vagina, from 73 percent to 98 percent; the penis, 76 percent to 90 percent; and the anus, 87 percent to 96 percent.

To compile these data, researchers examined 2,670 HPV DNA tissue samples from seven population-based <u>cancer</u> registries.

Study authors intend to perform additional research in the future to follow up on their estimate of how well the current vaccines protect against HPV-associated cancers.

More information: *Journal of the National Cancer Institute*: 2015 May: US Assessment of HPV Types in Cancers: Implications for Current and 9-Valent HPV Vaccines.

Provided by Cedars-Sinai Medical Center

Citation: 80 percent of cervical cancers found to be preventable with latest 9-valent HPV vaccine (2015, May 11) retrieved 3 May 2024 from https://medicalxpress.com/news/2015-05-percent-cervical-cancers-latest-valent.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.