

Population-based FIT screening can reduce CRC mortality

May 28 2015



(HealthDay)—Population-based fecal immunochemical testing (FIT) screening in adults aged 50 to 69 years can reduce colorectal cancer (CRC) mortality, according to a study published online May 20 in *Cancer*.

Han-Mo Chiu, Ph.D., from National Taiwan University, and colleagues conducted a prospective study of the follow-up of about five million Taiwanese from 2004 to 2009. CRC mortality was compared for a screened group and an unscreened group in a [population-based CRC screening](#) service targeting 50- to 69-year-olds. Overall, 21.4 percent of the 5,417,699 subjects aged 50 to 69 years in the population participated in the biennial screening program by 2009.

The researchers found that, with a maximum follow-up of six years, the actual effectiveness of FIT in reducing CRC mortality was 62 percent (relative rate for screened versus unscreened group, 0.38). After adjustment for self-selection bias, the 21.4 percent coverage of the population receiving FIT correlated with a significant 10 percent reduction in CRC [mortality](#) (relative rate, 0.9).

"Although such findings are informative for health decision makers, continued follow-up of this large cohort will be required to estimate the long-term impact of FIT screening if the covered population is expanded," the authors write.

More information: [Abstract](#)
[Full Text](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: Population-based FIT screening can reduce CRC mortality (2015, May 28) retrieved 4 May 2024 from <https://medicalxpress.com/news/2015-05-population-based-screening-crc-mortality.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
