

Aerobic fitness measures predict post-AAA complications

June 12 2015



(HealthDay)—Cardiopulmonary exercise testing variables can predict postoperative complications after abdominal aortic aneurysm repair, according to a study published in the June issue of *Anaesthesia*.

Hashem M. Barakat, M.B.B.S., from the University of Hull & Hull York Medical School in the United Kingdom, and colleagues examined whether measures of aerobic fitness can predict organ-specific complications after abdominal aortic aneurysm repair. Cardiopulmonary [exercise testing](#) data were collected prospectively over two years for 130 patients.

The researchers found that a decreased anaerobic threshold and open repair correlated with cardiac complications (odds ratios [ORs], 0.55 and 6.99, respectively). Increased ventilatory equivalent for carbon dioxide and open repair also correlated with pulmonary complications (ORs,

1.18 and 14.29, respectively). Hospital and critical care lengths of stay were shorter for patients who had an endovascular repair. There was no correlation for measures of fitness with 30-day mortality or length of hospital stay.

"Cardiopulmonary exercise testing variables, therefore, seem to predict different [postoperative complications](#) following [abdominal aortic aneurysm](#) repair, which adds value to their routine use in risk stratification and optimization of perioperative care," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: Aerobic fitness measures predict post-AAA complications (2015, June 12) retrieved 24 May 2024 from <https://medicalxpress.com/news/2015-06-aerobic-post-aaa-complications.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--