

Doctors worry about return of vaccinepreventable ills in kids

June 12 2015, by Amy Norton, Healthday Reporter



When immunization rates go down, 'herd' immunity also declines, experts note.

(HealthDay)—Although most U.S. children are getting their routine vaccinations, recent trends have experts concerned that Americans will lose some of the "herd immunity" that has long protected many from serious infections.

The vast majority of U.S. kids are up-to-date with routine jabs against once-common infections like polio, measles, mumps, whooping cough and chickenpox. But that "coverage" varies from state to state, according to the latest figures from the U.S. Centers for Disease Control and Prevention.

In 2013, there were 17 states where less than 90 percent of 1.5- to 3-yearolds had gotten their first dose of the measles-mumps-rubella (MMR) vaccine, the CDC found.



And nationally, recent years have seen a small dip in the percentage of young children who are up-to-date with some other vaccines, according to Dr. Mark Sawyer, an infectious disease specialist at Rady Children's Hospital and the University of California, San Diego, School of Medicine.

It's concerning, Sawyer said, because some parents are opting to skip or delay vaccinations not for medical reasons, but for "personal beliefs"—mainly, unsupported fears about <u>vaccine safety</u>.

Normally, in a community where most people are immunized, the unvaccinated few are still protected by the cocoon of immune folks around them. That so-called herd immunity shields infants who are too young for certain vaccines, and people who can't be immunized because of a medical condition.

"But herd immunity is only as good as the herd you're in," said Sawyer, who spoke this week at a news conference sponsored by the March of Dimes.

"I'm concerned that we're slowly losing the battle, and we're about to lose some of this herd immunity," Sawyer said. "How about Disneyland and the measles outbreak as an example?"

He was referring to the measles outbreak that, as of May 29, has sickened 173 people in 21 U.S. states and Washington, D.C., according to the CDC. The main outbreak was traced to Disneyland, in California.

The CDC says it probably began with an unvaccinated traveler who became infected in another country before visiting the amusement park, where he or she encountered other unvaccinated people.

"It's remarkable that this one case managed to spread to over 100



people," Sawyer said.

Some U.S. parents, he noted, think diseases like measles and whooping cough are relics of the past, so skipping or delaying their children's vaccinations will cause no harm. But unvaccinated, and unknowingly infected, travelers flow in and out of the United States all the time, Sawyer said—and those travelers are typically the "index case" in disease outbreaks here.

"They're not from third world countries, either," Sawyer noted. "They're often from highly developed European countries where <u>vaccination rates</u> are low and they've let the genie out of the bottle."

What's more, parents' worries about vaccine safety have no science behind them, emphasized both Sawyer and the CDC's Dr. Sonja Rasmussen, who also spoke at the conference.

The notion that the MMR vaccine causes autism has lingered for years, even though the 1998 study that proposed the theory was later found to be fraudulent.

Another misperception, Sawyer said, is that the current vaccine schedule subjects babies and preschoolers to too many shots in a short period. As a result, some parents ask their doctors to delay certain vaccines—and, recent research shows, they often get their wish.

However, Sawyer said, studies have found no evidence that the vaccine schedule "overwhelms" babies' immune systems.

In contrast, no studies have looked at whether delayed or "alternative" vaccination schedules are as effective as the recommended one.

"The recommended schedule is not put together casually," Sawyer said.



"There's a lot of science behind it."

Rasmussen underscored another strategy for protecting young infants: Vaccinating moms-to-be and family members.

Pregnant women should get a flu shot, she said, because it lowers their risk of a severe case of the flu, and the pregnancy complications that can cause. Plus, moms pass on some of those flu-fighting antibodies to their babies.

That's important, Rasmussen said, because infants cannot get the flu vaccine until the age of 6 months.

Pregnant women should also get the Tdap booster shot against whooping cough, because immunity to that infection wanes over time, Rasmussen said. The same goes for family members and anyone caring for the baby, since infants do not get their first shot against whooping cough until they're 2 months old.

For adults and older kids, <u>whooping cough</u> is miserable but relatively mild. That's not true for infants, Rasmussen pointed out.

"It's babies who are at risk of severe infections and death," she said.

Sawyer stressed the point of community protection. High vaccination rates keep vulnerable groups, like infants without full immunity, safe from serious infections, he said.

"When you decide not to vaccinate your child," Sawyer said, "it affects the whole community."

More information: The CDC has more on vaccine safety.



Copyright © 2015 <u>HealthDay</u>. All rights reserved.

Citation: Doctors worry about return of vaccine-preventable ills in kids (2015, June 12) retrieved 27 June 2024 from https://medicalxpress.com/news/2015-06-doctors-vaccine-preventable-ills-kids.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.