

Three Ebola virus variants identified in Guinea

June 24 2015



A map of Ebola virus variants circulating in Guinea and Sierra Leone. Credit: Creative Commons - *Nature*

Sequencing the genome of Ebola virus strains circulating in Guinea has



allowed scientists at the Institut Pasteur in Dakar and in Paris, the CNRS and the University of Sydney to retrace the spread of the virus and monitor its evolution in the country where the outbreak started. This research reveals the co-circulation in Guinea, particularly in the urban regions of the capital and neighboring towns, of three distinct variants of the virus whose mutations are described in an article published in *Nature*. Characterization of the genetic variations of the virus is crucial to ensure the continued efficacy of diagnostic tools and for the development of effective treatments and vaccines.

The Ebola epidemic has been ongoing in West Africa for over a year, with 27,341 reported cases, of which 11,184 have been fatal. The source of the epidemic has been traced to a forested area in South-East Guinea, from where it rapidly spread to the capital, Conakry, and to neighboring countries. In March 2014, the Institut Pasteur of Dakar set up a mobile laboratory at Donka hospital (Conakry), to provide diagnostic services throughout Guinea. The involvement of volunteers from the Institut Pasteur and its network, in Conakry and other regions of Guinea such as Macenta, was constant during the entire epidemic.

Monitoring the development of the Ebola <u>virus</u> genome is key for developing better treatment strategies, designing effective vaccines and ensuring the continued efficacy of <u>diagnostic tools</u>. To this end, scientists at the Institut Pasteur in Dakar and Paris, the CNRS and the University of Sydney contributed to the international efforts to combat the disease by characterizing the Ebola virus isolates circulating in Guinea between July and November 2014.

Sequencing of the virus from rare samples containing only small amounts of biological material was optimized in a collaborative effort with scientists from the Broad Institute (Cambridge, USA), who were already deployed in Sierra Leone. Analysis of these sequences reveals the existence of three distinct variants of the virus co-circulating in



Guinea, and a very different dynamic than that observed in Sierra Leone and Liberia.

The first variant (GUI-1 - shown in red on the map) is closely related to viruses sampled at the start of the epidemic in March 2014. This variant is found only in Guinea, both in urban areas (Conakry) and forested regions.

The second variant (GUI-2 - shown in blue on the map) is related to the viruses circulating in Sierra Leone, but could correspond to parallel evolution in Guinea. These sequences represent the missing link that led to two separate introductions of Ebola in Mali, in October and November 2014.

The third variant (SLE-GUI-3 - shown in orange on the map) was identified in Conakry and the surrounding towns (Forecariah, Dalaba and Coyah). The marked similarities between this variant and the viruses found in Sierra Leone, combined with epidemiological data, highlight multiple reintroductions of Ebola from Sierra Leone to the region around Conakry.

Each variant is defined by a combination of mutations affecting different viral proteins, in particular the VP35 protein, which may be a virulence factor, the virus envelope glycoprotein, which may alter the immune system's perception of the virus, or the polymerase, which is usually a more conserved viral region.

While this study highlights the genetic diversity of the viruses circulating in Guinea during the epidemic's spread, it also shows that the mutation rate is well within the margins previously described for this type of virus. Monitoring viral variations is a strong complement to epidemiological studies retracing transmission chains, and will enable better targeting of response strategies in the face of potential new epidemics. Lastly, studies



of the Ebola virus genetic variations, which have quickly been made available to the scientific community, will help optimize the treatments and vaccines under development.

More information: Distinct lineages of Ebola virus in Guinea during the 2014 West African epidemic, *Nature*, June 24, 2015.

Provided by Institut Pasteur

Citation: Three Ebola virus variants identified in Guinea (2015, June 24) retrieved 2 May 2024 from <u>https://medicalxpress.com/news/2015-06-ebola-virus-variants-guinea.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.